arXiv:1303.6449v1 [math.PR] 26 Mar 2013

Dirichlet Heat Kernel Estimates for Rotationally Symmetric Lévy processes

Zhen-Qing Chen[*] **Panki Kim**[†] and **Renming Song**[‡]

(March 26, 2013)

Abstract

In this paper, we consider a large class of purely discontinuous rotationally symmetric Lévy processes. We establish sharp two-sided estimates for the transition densities of such processes killed upon leaving an open set D. When D is a κ-fat open set, the sharp two-sided estimates are given in terms of surviving probabilities and the global transition density of the Lévy process. When D is a $C^{1,1}$ open set and the Lévy exponent of the process is given by $\Psi(\xi) = \phi(|\xi|^2)$ with ϕ being a complete Bernstein function satisfying a mild growth condition at infinity, our two-sided estimates are explicit in terms of Ψ, the distance function to the boundary of D and the jumping kernel of X, which give an affirmative answer to the conjecture posted in [16]. Our results are the first sharp two-sided Dirichlet heat kernel estimates for a large class of symmetric Lévy processes with general Lévy exponents. We also derive an explicit lower bound estimate for symmetric Lévy processes on $\mathbb{R}^d$ in terms of their Lévy exponents.

AMS 2000 Mathematics Subject Classification: Primary 60J35, 47G20, 60J75; Secondary 47D07

Keywords and phrases: Lévy processes, subordinate Brownian motion, heat kernel, transition density, Dirichlet transition density, Green function, exit time, Lévy system, boundary Harnack inequality, parabolic Harnack inequality

1 Introduction

Due to their importance in theory and applications, fine potential theoretical properties of Lévy processes have been under intense study recently. The transition density $p(t, x, y)$ of a Lévy process is the heat kernel of the generator of the process. However, the transition density (if it exists) of a general Lévy process rarely admits an explicit expression. Thus obtaining sharp estimates on $p(t, x, y)$ is a fundamental problem both in probability theory and in analysis.

---

[*] Research partially supported by NSF Grant DMS-1206276 and NNSFC Grant 11128101.

[†] This work was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) grant funded by the Korea government(MEST) (2012-0000940).

[‡] Research supported in part by a grant from the Simons Foundation (208236).

The generator of a discontinuous Lévy process is an integro-differential operator and so it is a non-local operator. Recently, quite a few people in PDE are interested in problems related to non-local operators; see, for example, [5, 6, 7, 24] and the references therein.

When X is a symmetric diffusion on $\mathbb{R}^d$ whose infinitesimal generator is a uniformly elliptic and bounded divergence form operator, it is well-known that $p(t, x, y)$ enjoys the celebrated Aronson's Gaussian type estimates. When X is a pure jump symmetric process on $\mathbb{R}^d$, sharp estimates on $p(t, x, y)$ have been studied in [8, 9, 10, 17, 18] recently, which can be viewed as the counterpart of Aronson's estimates for non-local operators.

Due to the complication near the boundary, two-sided estimates for the transition densities of discontinuous Lévy processes killed upon leaving an open set D (equivalently, the Dirichlet heat kernels) have been established very recently for a few particular processes only. The first of such estimates is obtained in [11], where we succeeded in establishing sharp two-sided estimates for the heat kernel of the fractional Laplacian $\Delta^{\alpha/2} := -(-\Delta)^{\alpha/2}$ with zero exterior condition on D^c (or equivalently, the transition density of the killed symmetric α-stable process) in any $C^{1,1}$ open set D. The approach developed in [11] provides a road map for establishing sharp two-sided heat kernel estimates of other jump processes in open subsets of $\mathbb{R}^d$. The ideas of [11] were adapted to establish sharp two-sided heat kernel estimates of relativistic stable processes and mixed stable processes in $C^{1,1}$ open subsets of $\mathbb{R}^d$ in [14, 13] respectively. In all these cases, the characteristic exponents of these Lévy processes admit explicit expressions, the boundary decay rates of the Dirichlet heat kernels are suitable powers of the distance to the boundary. On the other hand, a Varopoulos type two-sided Dirichlet heat kernel estimate of symmetric stable processes in κ-fat open sets was derived in [4]; this type of estimates is expressed in terms of surviving probabilities and the global transition density of the symmetric stable process.

The objective of this paper is to establish sharp two-sided estimates on the transition density $p_D(t, x, y)$ for a large class of purely discontinuous rotationally symmetric Lévy processes. Unlike the cases considered in [4, 11, 13, 14], the characteristic exponents of the symmetric Lévy processes considered in this paper are quite general, satisfying only certain mild growth condition at ∞. Moreover, the boundary decay rate of $p_D(t, x, y)$ is no longer some power of the distance to the boundary. The analysis of the precise boundary behavior of $p_D(t, x, y)$ is quite challenging and delicate. The main tools to obtain the precise boundary behavior of $p_D(t, x, y)$ are two versions of the boundary Harnack principle obtained in [19, 21]. In this paper we combine the approaches developed in [4, 11] with these boundary Harnack principles to obtain sharp two-sided estimates for $p_D(t, x, y)$, which cover the main results in [4, 11, 13, 14] and much more.

Suppose that $S = (S_t : t \geq 0)$ is a subordinator with Laplace exponent ϕ, that is, S is a nonnegative Lévy process with $S_0 = 0$ and $\mathbb{E}\left[e^{-\lambda S_t}\right] = e^{-t\phi(\lambda)}$ for every $t, \lambda > 0$. The function ϕ can be written in the form

$$\phi(\lambda) = b\lambda + \int_0^\infty (1 - e^{-\lambda t})\mu(dt), \tag{1.1}$$

where $b \geq 0$ and μ is a measure on $(0, \infty)$ satisfying $\int_0^\infty (1 \wedge t)\mu(dt) < \infty$. The constant b is called the drift of the subordinator and μ the Lévy measure of the subordinator (or of ϕ). The function ϕ is a Bernstein function, i.e., it is C^∞, positive and $(-1)^{n-1}D^n\phi \geq 0$ for all $n \geq 1$. In particular, since $\phi(0) = 0$ and $\phi'' \leq 0$, the Bernstein function ϕ has the property that

$$\phi(\lambda r) \leq \lambda\phi(r) \qquad \text{for all } \lambda \geq 1 \text{ and } r > 0. \tag{1.2}$$

The Laplace exponent ϕ is said to be a complete Bernstein function if the Lévy measure μ of ϕ has a completely monotone density $\mu(t)$, i.e., $(-1)^n D^n \mu \geq 0$ for every non-negative integer n. For basic results on complete Bernstein functions, we refer the reader to [23].

Throughout this paper, we assume that ϕ is a complete Bernstein function satisfying the following growth condition at infinity (see [27]):

(A): There exist constants $\delta_1, \delta_2 \in (0,1)$, $a_1 \in (0,1)$, $a_2 \in (1,\infty)$ and $R_0 > 0$ such that

$$a_1 \lambda^{\delta_1} \phi(r) \leq \phi(\lambda r) \leq a_2 \lambda^{\delta_2} \phi(r) \qquad \text{for } \lambda \geq 1 \text{ and } r \geq R_0.$$

Note that it follows from the upper bound condition in **(A)** that ϕ has no drift.

Let $W = (W_t : t \geq 0)$ be a Brownian motion in $\mathbb{R}^d$ independent of the subordinator S. The subordinate Brownian motion $Y = (Y_t : t \geq 0)$ is defined by $Y_t := W_{S_t}$, which is a rotationally symmetric Lévy process with Lévy exponent $\phi(|\xi|^2)$. The infinitesimal generator of Y is $\mathcal{L}^Y := -\phi(-\Delta)$. Here and below for a function ψ on $[0,\infty)$, $\psi(-\Delta)$ is defined as a pseudo differential operator in terms of Fourier transform; that is, $\widehat{\psi(-\Delta)f}(\xi) := -\psi(|\xi|^2)\widehat{f}(\xi)$, where $\widehat{f}$ is the Fourier transform of a function f on $\mathbb{R}^d$. It is known that the Lévy measure of the process Y has a density given by $J(x) = j(|x|)$ where

$$j(r) := \int_0^\infty (4\pi t)^{-d/2} e^{-r^2/(4t)} \mu(t)\,dt, \qquad r > 0. \tag{1.3}$$

Note that the function $r \mapsto j(r)$ is continuous and decreasing on $(0,\infty)$.

We will assume that X is a purely discontinuous rotationally symmetric Lévy process with Lévy exponent $\Psi(\xi)$. Because of rotational symmetry, the function Ψ depends on $|\xi|$ only, and by a slight abuse of notation we write $\Psi(\xi) = \Psi(|\xi|)$. The infinitesimal generator of X is $\mathcal{L}^X := -\Psi(\sqrt{-\Delta})$. We further assume that the Lévy measure of X has a density with respect to the Lebesgue measure on $\mathbb{R}^d$, which is denoted by $J_X(x,y) = J_X(x-y) = j_X(|y-x|)$. That is,

$$\mathbb{E}_x\left[e^{i\xi\cdot(X_t - X_0)}\right] = e^{-t\Psi(|\xi|)} \qquad \text{for every } x \in \mathbb{R}^d \text{ and } \xi \in \mathbb{R}^d,$$

with

$$\Psi(|\xi|) = \int_{\mathbb{R}^d} (1 - \cos(\xi \cdot x)) J_X(x)\,dx. \tag{1.4}$$

We assume that $j_X(r)$ is continuous on $(0,\infty)$ and that there is a constant $\gamma > 1$ such that

$$\gamma^{-1} j(r) \leq J_X(r) \leq \gamma j(r) \quad \text{for all } r > 0. \tag{1.5}$$

This implies that J and J_X are comparable. Clearly (1.5) also implies that

$$\gamma^{-1}\phi(|\xi|^2) \leq \Psi(|\xi|) \leq \gamma\phi(|\xi|^2) \quad \text{for all } \xi \in \mathbb{R}^d. \tag{1.6}$$

We remark that under the above assumptions, X does not need to be a subordinate Brownian motion because j_X does not need to be monotone. For example, choose $\varepsilon > 0$ such that $2^{-1}j(1) < j(1+\varepsilon)$ and a continuous function h with $h(1) = \frac{1}{2}j(r)$, $h(1+\varepsilon) = 0$ and $0 \leq h(r) \leq \frac{1}{2}j(r)$ for all $r > 0$. Then $j_X(r) := j(r) - h(r)$ is not monotone and its corresponding Lévy process X (through Lévy exponent (1.4)) is not a subordinate Brownian motion.

Under the above setup, X has a continuous transition density $p(t,x,y)$ with respect to the Lebesgue measure on $\mathbb{R}^d$ (see [9]). Clearly, $p(t,x,y)$ is a function depending only on t and $|x-y|$, and so, by an abuse of notation, we also denote $p(t,x,y)$ by $p(t,|x-y|)$. For every open subset $D \subset \mathbb{R}^d$, we denote by X^D the subprocess of X killed upon exiting D. It is known (see [9]) that X^D has a transition density $p_D(t,x,y)$, with respect to the Lebesgue measure, which is jointly locally Hölder continuous. Note that $p_D(t,x,y)$ is the fundamental solution for $\mathcal{L}^X = -\Psi(\sqrt{-\Delta})$ in D with zero exterior condition and so it can also be called the Dirichlet heat kernel of $\mathcal{L}^X$ in D. The purpose of this paper is to establish sharp two-sided estimates on $p_D(t,x,y)$. The following two conditions will be needed for some of the results in this paper when D is unbounded.

(B): There exist constants $C_1 > 0$ and $C_2 \in (0,1]$ such that

$$p(t,u) \le C_1 p(t, C_2 r) \qquad \text{for } t \in (0,1] \text{ and } u \ge r > 0.$$

(C): There exist constants $C_3 > 0$ and $C_4 \in (0,1]$ such that

$$p(t,r) \le C_3 t j(C_4 r) \qquad \text{for } t \in (0,1] \text{ and } r > 0.$$

Throughout this paper we will use Φ to denote the function

$$\Phi(r) = \frac{1}{\phi(r^{-2})}, \qquad r > 0. \tag{1.7}$$

Note that in particular it follows from (1.2) that

$$\Phi(2r) = \frac{1}{\phi(r^{-2}/4)} \le \frac{1}{\phi^{-1}(r^{-2})/4} = 4\Phi(r) \qquad \text{for every } r > 0. \tag{1.8}$$

The inverse function of Φ will be denoted by the usual notation $\Phi^{-1}(r)$. Here and in the following, for $a,b \in \mathbb{R}$, $a \wedge b := \min\{a,b\}$ and $a \vee b := \max\{a,b\}$.

Remark 1.1 (i) The condition **(B)** is pretty mild. When X is a rotationally symmetric Lévy process such that $r \mapsto j_X(r)$ is decreasing, condition **(B)** holds for all $t > 0$ (see [25, Proposition]). In particular it holds for all subordinate Brownian motions with $C_1 = C_2 = 1$. In this special case, we can also see this using the following elementary argument: When X is a subordinate Brownian motion, $p(t,r) = \mathbb{E}[p_0(S_t, r)]$, where $p_0(t,|x-y|)$ is the transition density of the Brownian motion W. It follows immediately that $p(t,r)$ is decreasing in r and so **(B)** holds with $C_1 = C_2 = 1$.

(ii) Under condition **(A)**, condition **(B)** is weaker than **(C)**. Under condition **(A)**, we will show in this paper that there exists $c > 1$ such that

$$c^{-1}\big(\Phi^{-1}(t)^d \wedge t j(r)\big) \le p(t,r) \le c(\Phi^{-1}(t))^d \qquad \text{for } t \in (0,1] \text{ and } r > 0$$

(see Proposition 2.2 and Theorem 3.7 below). Thus condition **(C)** amounts to say that there exist constants $c \ge 1$ and $C_4 \in (0,1]$ such that for $t \in (0,1]$ and $r > 0$,

$$c^{-1}\big(\Phi^{-1}(t)^d \wedge t j(r)\big) \le p(t,r) \le c\big(\Phi^{-1}(t)^d \wedge t j(C_4 r)\big). \tag{1.9}$$

Since $j(r)$ is a decreasing function in r, (1.9) implies that condition **(B)** holds with $C_1 = c^2$ and $C_2 = C_4$.

(iii) Assume that condition **(A)** holds. It follows from [9, 10, 18] that, for every $R > 0$, there is a constant $c = c(T, R, \gamma, \phi) > 1$ so that (1.9) holds for $(t, r) \in (0, 1] \times (0, R]$ with $C_4 = 1$. (See Proposition 2.2 below.) So the assertions in conditions **(B)** and **(C)** are always satisfied for $0 < r \le u \le R$.

(iv) By [10, 18],

$$c^{-1}\left((\Phi^{-1}(t))^{-d} \wedge tj(C_4^{-1}r)\right) \le p(t,r) \le c\left((\Phi^{-1}(t))^{-d} \wedge tj(C_4 r)\right) \quad \text{for } (t,r) \in (0,1]\times(0,\infty),$$

and consequently conditions **(B)** and **(C)**, hold for a large class of discontinuous processes including mixed stable-like processes (with $C_4 = 1$) and relativistic stable-like processes (with $C_4 = 1$, see [14, Theorem 4.1]). □

Before stating the main results of this paper, we need first to set up some notations. Let $d \ge 1$. We denote the Euclidean distance between x and y in $\mathbb{R}^d$ by $|x-y|$ and denote by $B(x,r)$ the open ball centered at $x \in \mathbb{R}^d$ with radius $r > 0$; for any two positive functions f and g, $f \asymp g$ means that there is a positive constant $c \ge 1$ so that $c^{-1}g \le f \le c\,g$ on their common domain of definition; for any open $D \subset \mathbb{R}^d$ and $x \in D$, diam(D) stands for the diameter of D and $\delta_D(x)$ stands for the Euclidean distance between x and D^c.

Definition 1.2 *Let $0 < \kappa \le 1$. We say that a open set D is κ-fat if there is $R_1 > 0$ such that for all $x \in \overline{D}$ and all $r \in (0, R_1]$, there is a ball $B(A_r(x), \kappa r) \subset D \cap B(x,r)$. The pair (R_1, κ) is called the characteristics of the κ-fat open set D.*

The following factorization of the Dirichlet heat kernel is the first main result of this paper. Recall that C_1 and C_2 are the constants in condition **(B)**.

Theorem 1.3 *Let X be a purely discontinuous rotationally symmetric Lévy process with Lévy exponent Ψ and Lévy density J_X satisfying* (1.6) *and* (1.5) *respectively, where the complete Bernstein function ϕ satisfies* **(A)**. *Suppose that D is a κ-fat open set with characteristics (R_1, κ).*

(i) *For every $T > 0$, there exists $c_1 = c_1(R_1, \kappa, \gamma, T, d, \phi) > 0$ such that for $0 < t \le T$, $x, y \in D$,*

$$p_D(t,x,y) \ge c_1 \mathbb{P}_x(\tau_D > t)\mathbb{P}_y(\tau_D > t)\left(\Phi^{-1}(t)^{-d} \wedge tJ(x,y)\right). \tag{1.10}$$

(ii) *If D is unbounded, we assume in addition that condition* **(B)** *holds. For every $T > 0$, there exists $c_2 = c_2(C_1, C_2, R_1, \kappa, T, d, \gamma, \phi) > 0$ such that for $0 < t \le T$, $x, y \in D$,*

$$p_D(t,x,y) \le c_2 \mathbb{P}_x(\tau_D > t)\mathbb{P}_y(\tau_D > t)p(t, C_5 x, C_5 y), \tag{1.11}$$

where $C_5 = C_2^2/4$.

(iii) *Suppose in addition that D is bounded. Then there exists $c_3 = c_3(diam(D), R_1, \kappa, d, \gamma, \phi) > 1$ so that for all $(t,x,y) \in [3,\infty) \times D \times D$,*

$$c_3^{-1}\,\mathbb{P}_x(\tau_D > 1)\,\mathbb{P}_y(\tau_D > 1)\,e^{-t\lambda_1} \;\le\; p_D(t,x,y) \;\le\; c_3\,\mathbb{P}_x(\tau_D > 1)\,\mathbb{P}_y(\tau_D > 1)\,e^{-t\lambda_1},$$

where $-\lambda_1 < 0$ is the largest eigenvalue of the generator of X^D.

When X is a rotationally symmetric α-stable process in $\mathbb{R}^d$, that is, when $\Psi(\xi) = |\xi|^\alpha$ for some $\alpha \in (0, 2)$, parts (i) and (ii) of Theorem 1.3 are proved in [4].

Recall that C_3 and C_4 are the constants in condition **(C)**. Combining Theorem 1.3(i)–(ii) and Remark 1.1(ii), we have the following corollary.

Corollary 1.4 *Let X be a purely discontinuous rotationally symmetric Lévy process with Lévy exponent Ψ and Lévy density J_X satisfying* (1.6) *and* (1.5) *respectively, where the complete Bernstein function ϕ satisfies* **(A)**. *Suppose that D is a κ-fat open set with characteristics (R_1, κ). If D is unbounded, we assume in addition that condition* **(C)** *holds. For every $T > 0$, there exist $c_1 = c_1(R_1, \kappa, T, d, \gamma, \phi) > 0$ and $c_2 = c_2(C_3, C_4, R_1, \kappa, T, d, \gamma, \phi) > 0$ such that for $0 < t \le T$, $x, y \in D$,*

$$\begin{aligned} &c_1 \mathbb{P}_x(\tau_D > t)\mathbb{P}_y(\tau_D > t) \left(\Phi^{-1}(t)^{-d} \wedge t j(|x-y|) \right) \\ \le\, &p_D(t, x, y) \le c_2 \mathbb{P}_x(\tau_D > t)\mathbb{P}_y(\tau_D > t) \left(\Phi^{-1}(t)^{-d} \wedge t j(C_6|x-y|) \right), \end{aligned}$$

where $C_6 = C_4^3/4$.

The second main result of this paper is on explicit sharp Dirichlet heat kernel estimates for subordinate Brownian motions in $C^{1,1}$ open sets. So in the remainder of this section, we assume that $X = Y$, a subordinate Brownian motion with Lévy exponent $\Psi(\xi) = \phi(|\xi|^2)$.

Recall that an open set D in $\mathbb{R}^d$ (when $d \ge 2$) is said to be a (uniform) $C^{1,1}$ open set if there exist a localization radius $R_2 > 0$ and a constant $\Lambda > 0$ such that for every $z \in \partial D$, there exist a $C^{1,1}$-function $\psi = \psi_z : \mathbb{R}^{d-1} \to \mathbb{R}$ satisfying $\psi(0) = 0$, $\nabla\psi(0) = (0, \dots, 0)$, $\|\nabla\psi\|_\infty \le \Lambda$, $|\nabla\psi(x) - \nabla\psi(z)| \le \Lambda|x - z|$, and an orthonormal coordinate system CS_z with its origin at z such that

$$B(z, R_2) \cap D = \{y = (\widetilde{y}, y_d) \text{ in } CS_z : |y| < R_2, y_d > \psi(\widetilde{y})\}.$$

The pair (R_2, Λ) is called the characteristics of the $C^{1,1}$ open set D. Note that a $C^{1,1}$ open set D with characteristics (R_2, Λ) can be unbounded and disconnected; the distance between two distinct components of D is at least R_2. By a $C^{1,1}$ open set in $\mathbb{R}$ we mean an open set which can be written as the union of disjoint intervals so that the minimum of the lengths of all these intervals is positive and the minimum of the distances between these intervals is positive.

Here is the second main result of this paper, which gives an affirmative answer to the Conjecture posed in [16]. In view of Remark 1.1, it extends the main results of [11, 13, 14]. Recall that condition **(B)** holds with $C_1 = C_2 = 1$ for any subordinate Brownian motion.

Theorem 1.5 *Suppose that X is a subordinate Brownian motion with Lévy exponent $\Psi(\xi) = \phi(|\xi|^2)$ with ϕ being a complete Bernstein function satisfying condition* **(A)**. *Let D be a $C^{1,1}$ open subset of $\mathbb{R}^d$ with characteristics (R_2, Λ).*

(i) *For every $T > 0$, there exists $c_1 = c_1(R_2, \Lambda, T, d, \phi) > 0$ such that for all $(t, x, y) \in (0, T] \times D \times D$,*

$$p_D(t, x, y) \ge c_1 \left(1 \wedge \frac{\Phi(\delta_D(x))}{t}\right)^{1/2} \left(1 \wedge \frac{\Phi(\delta_D(y))}{t}\right)^{1/2} \left(\Phi^{-1}(t)^{-d} \wedge t J(x, y)\right).$$

(ii) *For every $T > 0$, there exists $c_2 = c_2(R_2, \Lambda, T, d, \phi) > 0$ such that for all $(t, x, y) \in (0, T] \times D \times D$,*

$$p_D(t, x, y) \le c_2 \left(1 \wedge \frac{\Phi(\delta_D(x))}{t}\right)^{1/2} \left(1 \wedge \frac{\Phi(\delta_D(y))}{t}\right)^{1/2} p(t, |x - y|/4).$$

(iii) *Suppose in addition that D is bounded. For every $T > 0$, there exists $c_3 \ge 1$ depending only on diam(D), $\lambda, R_2, \Lambda, d, \phi$ and T so that for all $(t, x, y) \in [T, \infty) \times D \times D$,*

$$c_3^{-1}\, e^{-\lambda_1 t} \sqrt{\Phi(\delta_D(x))}\sqrt{\Phi(\delta_D(y))} \;\le\; p_D(t, x, y) \;\le\; c_3\, e^{-\lambda_1 t}\, \sqrt{\Phi(\delta_D(x))}\sqrt{\Phi(\delta_D(y))},$$

where $-\lambda_1 < 0$ is the largest eigenvalue of the generator of X^D.

Recall that C_3 and C_4 are the constants in condition **(C)**. Combining Theorem 1.5(i)–(ii) and Remark 1.1(ii), we have the following corollary.

Corollary 1.6 *Suppose that X is a subordinate Brownian motion with Lévy exponent $\Psi(\xi) = \phi(|\xi|^2)$ with ϕ being a complete Bernstein function satisfying condition* **(A)**. *Let D be a $C^{1,1}$ open subset of $\mathbb{R}^d$ with characteristics (R_2, Λ). If D is unbounded, we assume in addition that condition* **(C)** *holds. For every $T > 0$, there exist $c_1 = c_1(R_2, \Lambda, T, d, \phi) > 0$ and $c_2 = c_2(C_3, C_4, R_2, \Lambda, T, d, \phi) > 0$ such that for $0 < t \le T$, $x, y \in D$,*

$$\begin{aligned}
&c_1 \left(1 \wedge \frac{\Phi(\delta_D(x))}{t}\right)^{1/2} \left(1 \wedge \frac{\Phi(\delta_D(y))}{t}\right)^{1/2} \left(\Phi^{-1}(t)^{-d} \wedge tj(|x - y|)\right) \\
\le\, &p_D(t, x, y) \le c_2 \left(1 \wedge \frac{\Phi(\delta_D(x))}{t}\right)^{1/2} \left(1 \wedge \frac{\Phi(\delta_D(y))}{t}\right)^{1/2} \left(\Phi^{-1}(t)^{-d} \wedge tj(C_4|x - y|/4)\right).
\end{aligned}$$

When X is a rotationally symmetric α-stable process in $\mathbb{R}^d$, Theorem 1.5 is first established in [11]. Sharp two-sided Dirichlet heat kernel estimates in $C^{1,1}$ open sets are subsequently established in [12, 14, 13, 15] for censored stable processes, relativistic stable processes, mixed stable processes, and mixed Brownian motion and stable processes, respectively. By integrating the two-sided heat kernel estimates in Theorem 1.5 with respect to t, we obtain the two-sided estimates on the Green function $G_D(x, y) := \int_0^\infty p_D(t, x, y)dt$ (see Theorem 7.3 below), which extend [20, Theorem 1.1].

Condition **(A)** is a very weak condition on the behavior of ϕ near infinity. Using the tables at the end of [23], one can come up plenty of explicit examples of complete Bernstein functions satisfying condition **(A)**. Here are a few of them.

(1) $\phi(\lambda) = \lambda^{\alpha/2}$, $\alpha \in (0, 2]$ (symmetric α-stable process);
(2) $\phi(\lambda) = (\lambda + m^{2/\alpha})^{\alpha/2} - m$, $\alpha \in (0, 2)$ and $m > 0$ (relativistic α-stable process);
(3) $\phi(\lambda) = \lambda^{\alpha/2} + \lambda^{\beta/2}$, $0 < \beta < \alpha < 2$ (mixed symmetric α- and β-stable processes);
(4) $\phi(\lambda) = \lambda^{\alpha/2}(\log(1 + \lambda))^p$, $\alpha \in (0, 2)$, $p \in [-\alpha/2, (2 - \alpha)/2]$.

Now we give a way of constructing less explicit complete Bernstein functions that have very general asymptotic behavior at infinity. Suppose that $\alpha \in (0, 2)$ and ℓ is a positive function on $(0, \infty)$ which is slowly varying at infinity. We further assume that $t \to t^{\alpha/2}\ell(t)$ is a right continuous increasing function with $\lim_{t\to 0} t^{\alpha/2}\ell(t) = 0$ (so $\int_0^\infty (1 + t)^{-2} t^{\alpha/2}\ell(t)dt < \infty$). Then the function

$$f(\lambda) := \int_0^\infty (\lambda + t)^{-2} t^{\alpha/2}\ell(t)dt$$

is a Stieltjes function, and so the function

$$\phi(\lambda) := \frac{1}{f(\lambda)} = \left(\int_0^\infty (\lambda+t)^{-2} t^{\alpha/2} \ell(t) dt \right)^{-1}$$

is a complete Bernstein function (see [23, Theorem 7.3]). It follows from [26, Lemma 6.2] that $\phi(\lambda) \asymp \lambda^{\alpha/2}\ell(\lambda)$ when $\lambda \geq 2$.

The rest of the paper is organized as follows. Section 2 recalls and collects some preliminary results that will be used in the sequel, including on-diagonal heat kernel estimates and the boundary Harnack principle. Section 3 presents the interior lower bound heat kernel estimates, including an explicit lower bound estimate for symmetric Lévy processes on $\mathbb{R}^d$. The proof of the short time factorization result for $p_D(t,x,y)$ (that is, Theorem 1.3(i) and (ii)) is given in Section 4, while the proof of Theorem 1.5(i) and (ii) is given in Section 5. The large time heat kernel estimates are proved in Section 6. The Green function estimates for subordinate Brownian motions in bounded $C^{1,1}$ open sets are derived in Section 7 from the two-sided Dirichlet heat kernel estimates in Theorem 1.5. The derivation, however, requires quite some effort.

Throughout this paper, $d \geq 1$ and the constants R_0, R_1, R_2, Λ, κ, δ_1, δ_2, C_1, C_2, C_3, C_4, C_5 and C_6 will be fixed. We use $c_1, c_2, \cdots$ to denote generic constants, whose exact values are not important and can change from one appearance to another. The labeling of the constants $c_1, c_2, \cdots$ starts anew in the statement of each result. The dependence of the constant c on the dimension d will not be mentioned explicitly. We will use ":=" to denote a definition, which is read as "is defined to be". We will use ∂ to denote a cemetery point and for every function f, we extend its definition to ∂ by setting $f(\partial) = 0$. We will use dx to denote the Lebesgue measure in $\mathbb{R}^d$. For a Borel set $A \subset \mathbb{R}^d$, we also use $|A|$ to denote its Lebesgue measure.

2 Preliminary

In the first part of this section, we assume that X is a purely discontinuous rotationally symmetric Lévy process with Lévy exponent Ψ and Lévy density J_X satisfying (1.6) and (1.5) respectively, where the complete Bernstein function ϕ satisfies **(A)**.

A function $u : \mathbb{R}^d \mapsto [0,\infty)$ is said to be harmonic in an open set $D \subset \mathbb{R}^d$ with respect to X if for every open set B whose closure is a compact subset of D,

$$u(x) = \mathbb{E}_x \left[u(X_{\tau_B}) \right] \qquad \text{for every } x \in B. \tag{2.1}$$

A function $u : \mathbb{R}^d \mapsto [0,\infty)$ is said to be regular harmonic in an open set $D \subset \mathbb{R}^d$ with respect to X if

$$u(x) = \mathbb{E}_x \left[u(X_{\tau_D}) \right] \qquad \text{for every } x \in D.$$

Clearly, a regular harmonic function in D is harmonic in D.

Very recently the following form of the boundary Harnack principle is established in [21].

Theorem 2.1 ([21, Theorem 1.1(i)]) *There exists $c = c(\phi, \gamma) > 0$ such that for any $z_0 \in \mathbb{R}^d$, any open set $D \subset \mathbb{R}^d$, any $r \in (0,1)$ and any nonnegative functions u, v in $\mathbb{R}^d$ which are regular harmonic in $D \cap B(z_0, r)$ with respect to X and vanish in $D^c \cap B(z_0, r)$, we have*

$$\frac{u(x)}{v(x)} \leq c \frac{u(y)}{v(y)} \qquad \textit{for all } x, y \in D \cap B(z_0, r/2).$$

Recall also from [21] that j enjoys the following properties: for every $R>0$,

$$j(r) \asymp \frac{1}{r^d\,\Phi(r)} \quad \text{for } r\in(0,R], \tag{2.2}$$

and there exists $c>0$ such that

$$j(r) \le cj(r+1) \qquad \text{for } r\ge 1. \tag{2.3}$$

It follows from (1.8) and (2.2) that both the function Φ defined by (1.7) and the function j satisfy a doubling property; that is, for every constant $R>0$, there is a constant $c>1$ so that

$$\Phi(2r)\le 4\,\Phi(r) \quad \text{and} \quad j(r)\le cj(2r) \quad \text{for every } r\in(0,R]. \tag{2.4}$$

Moreover, under condition **(A)**, for any given $R>0$, $j(r)$ satisfies all the conditions in [18] for $r\in(0,R)$. Thus we have the following two-sided estimates for $p(t,x,y)$ from [9, 10]:

Proposition 2.2 *For any $T>0$, there exists $c_1=c_1(T,R,\gamma,\phi)>0$ such that*

$$p(t,x,y)\le c_1\,(\Phi^{-1}(t))^{-d} \qquad \text{for } (t,x,y)\in(0,T]\times\mathbb{R}^d\times\mathbb{R}^d. \tag{2.5}$$

For any $T,R>0$, there exists $c_2=c_2(T,R,\gamma,\phi)>1$ such that for all $(t,x,y)\in[0,T]\times\mathbb{R}^d\times\mathbb{R}^d$ with $|x-y|<R$,

$$c_2^{-1}\left((\Phi^{-1}(t))^{-d}\wedge tJ(x,y)\right)\le p(t,x,y)\le c_2\left((\Phi^{-1}(t))^{-d}\wedge tJ(x,y)\right). \tag{2.6}$$

Proof. (2.5) is given in the first display on page 1073 of [9]. By [9, Theorem 2.4] and (1.5), there exist $T_*>0$ and R_* such that for all $(t,x,y)\in[0,T_*]\times\mathbb{R}^d\times\mathbb{R}^d$ with $|x-y|<R_*$, (2.6) holds. Now we assume $R>R_*$. We construct Z from X by removing jumps of size larger than R via Meyer's construction (see [22]). Let $p_Z(t,x,y)$ be the transition density of Z. By [1, Lemma 3.6] and [2, Lemma 3.1(c)] we have for every $t>0$ and $x,y\in\mathbb{R}^d$,

$$e^{-t\|\mathcal{J}_R\|_\infty}p_Z(t,x,y)\ \le\ p(t,x,y)\ \le\ p_Z(t,x,y)+t\|J_R\|_\infty, \tag{2.7}$$

where

$$J_R(x,y):=J_X(x,y)\mathbf{1}_{\{|x-y|>R\}} \quad \text{and} \quad \mathcal{J}_R(x):=\int_{\mathbb{R}^d}J_R(x,y)dy. \tag{2.8}$$

Applying [10, Theorem 1.4] and its proof to (2.7), we get

$$c_1e^{-t\|\mathcal{J}_R\|_\infty}\left((\Phi^{-1}(t))^{-d}\wedge tJ_X(x,y)\right)\ \le\ p(t,x,y)\ \le\ c_2\left((\Phi^{-1}(t))^{-d}\wedge tJ_X(x,y)\right)+t\|J_R\|_\infty. \tag{2.9}$$

(2.6) now follows from (1.5). □

The function $J_X(x,y)$ gives rise to a Lévy system for X, which describes the jumps of the process X: for any non-negative measurable function f on $\mathbb{R}_+\times\mathbb{R}^d\times\mathbb{R}^d$ with $f(s,y,y)=0$ for all $y\in\mathbb{R}^d$ and stopping time T (with respect to the filtration of X),

$$\mathbb{E}_x\left[\sum_{s\le T}f(s,X_{s-},X_s)\right]=\mathbb{E}_x\left[\int_0^T\left(\int_{\mathbb{R}^d}f(s,X_s,y)J_X(X_s,y)dy\right)ds\right]. \tag{2.10}$$

(See, for example, [17, Proof of Lemma 4.7] and [18, Appendix A].)

When $\Psi(|\xi|) = \phi(|\xi|^2)$ and ϕ is a complete Bernstein function satisfying condition **(A)**, the following boundary Harnack principle on a $C^{1,1}$ open subset with explicit decay rate is established in [19] (see also [20]).

Theorem 2.3 ([19, Theorem 1.5]) *Suppose that X is a rotationally symmetric Lévy process with Lévy exponent $\Psi(\xi) = \phi(|\xi|^2)$ with ϕ being a complete Bernstein function satisfying condition* **(A)**. *Assume that D is a (possibly unbounded) $C^{1,1}$ open set in $\mathbb{R}^d$ with characteristics (R_2, Λ). Then there exists $c = c(R_2, \Lambda, \phi, d) > 0$ such that for $r \in (0, (R_2 \wedge 1)/4]$, $Q \in \partial D$ and any nonnegative function u in $\mathbb{R}^d$ that is harmonic in $D \cap B(Q, r)$ with respect to X and vanishes continuously on $D^c \cap B(Q, r)$, we have*

$$\frac{u(x)}{\Phi(\delta_D(x))^{1/2}} \le c\,\frac{u(y)}{\Phi(\delta_D(y))^{1/2}} \qquad \text{for every } x, y \in D \cap B(Q, r/2). \tag{2.11}$$

3 Interior lower bound estimate

In this section, we assume that X is a purely discontinuous rotationally symmetric Lévy process with Lévy exponent Ψ and Lévy density J_X satisfying (1.6) and (1.5) respectively, where the complete Bernstein function ϕ satisfies **(A)**. We will give some preliminary lower bounds on $p_D(t, x, y)$ and $p(t, x, y)$. We first recall the following from [9, 18]. Recall that $\mathcal{J}_R$ was defined in (2.8).

Lemma 3.1 *For any positive constant M, there exist $\varepsilon = \varepsilon(M, \gamma, \phi) > 0$ and $c = c(M, \gamma, \phi) \ge 1$ such that for all $r \in (0, M]$ and $z \in \mathbb{R}^d$,*

$$\mathbb{P}_z\left(\tau_{B(z,r)} > \varepsilon\Phi(r)\right) \ge 2^{-1} e^{-\|\mathcal{J}_1\|_\infty} \tag{3.1}$$

and

$$c^{-1}\,\Phi(r) \;\le\; \mathbb{E}_z\left[\tau_{B(z,r)}\right] \;\le\; c\,\Phi(r). \tag{3.2}$$

Proof. (3.1) follows directly from [9, Lemma 2.5]. We then have

$$\mathbb{E}_z\left[\tau_{B(z,r)}\right] \ge \varepsilon\Phi(r)\mathbb{P}_z(\tau_{B(z,r)} > \varepsilon\,\Phi(r)) > 2^{-1}e^{-\|\mathcal{J}_1\|_\infty}\varepsilon\Phi(r).$$

On the other hand, by the Lévy system for X in (2.10), (1.5), (2.2) and (2.4), we have

$$\begin{aligned} 1 \;&\ge\; \mathbb{P}_z\left(X_{\tau_{B(z,r)}} \in B(z,2r)^c\right) = \mathbb{E}_z\left[\int_0^{\tau_{B(z,r)}} \int_{B(z,2r)^c} J_X(X_s, y)dyds\right] \\ &\ge\; c_1\mathbb{E}_z\left[\int_0^{\tau_{B(z,r)}} \int_{B(z,3r)\setminus B(z,2r)} J(X_s, y)dyds\right] \ge c_2\frac{|B(z,3r)\setminus B(z,2r)|}{r^d\Phi(r)}\,\mathbb{E}_z\left[\tau_{B(z,r)}\right], \end{aligned}$$

which yields $\mathbb{E}_z\left[\tau_{B(z,r)}\right] \le c\Phi(r)$. □

Lemma 3.2 *For any positive constants a and R, there exists $c = c(a, R, \gamma, \phi) > 0$ such that for all $z \in \mathbb{R}^d$ and $r \in (0, R]$,*

$$\inf_{y \in B(z,r/2)} \mathbb{P}_y\left(\tau_{B(z,r)} > a\Phi(r)\right) \;\ge\; c.$$

Proof. By (3.1) and (2.4), there exists $\varepsilon_1 = \varepsilon_1(R, \gamma, \phi) > 0$ such that for all $r \in (0, R]$,

$$\inf_{z\in\mathbb{R}^d} \mathbb{P}_z(\tau_{B(z,r/2)} > \varepsilon_1 \Phi(r)) \geq 2^{-1}e^{-\|\mathcal{J}_1\|_\infty}.$$

Thus it suffices to prove the lemma for $a > \varepsilon_1$. Applying the parabolic Harnack inequality [9, Theorem 5.2] at most $2 + [a/\varepsilon_1]$ times, we conclude that there exists $c_1 = c_1(a, R, \gamma, \phi) > 0$ such that for every $w, y \in B(z, r/2)$,

$$c_1\, p_{B(z,r)}(\varepsilon_1 \Phi(r), z, w) \;\leq\; p_{B(z,r)}(a\Phi(r), y, w).$$

Thus

$$\begin{aligned}
\mathbb{P}_y\left(\tau_{B(z,r)} > a\Phi(r)\right) &= \int_{B(z,r)} p_{B(z,r)}(a\Phi(r), y, w)dw \\
&\geq \int_{B(z,r/2)} p_{B(z,r)}(a\Phi(r), y, w)dw \\
&\geq c_1 \int_{B(z,r/2)} p_{B(z,r/2)}(\varepsilon_1\Phi(r), z, w)dw \\
&= c_1 \mathbb{P}_z(\tau_{B(z,r/2)} > \varepsilon_1\Phi(r)) \;\geq\; 2^{-1}e^{-\|\mathcal{J}_1\|_\infty}.
\end{aligned}$$

This proves the lemma. □

For the next four results, D is an arbitrary nonempty open set and we use the convention that $\delta_D(\cdot) \equiv \infty$ when $D = \mathbb{R}^d$.

Proposition 3.3 *Let $T > 0$ and $a > 0$ be constants. There exists $c = c(T, a, \gamma, \phi) > 0$ such that*

$$p_D(t, x, y) \;\geq\; c\,(\Phi^{-1}(t))^{-d} \tag{3.3}$$

for every $(t, x, y) \in (0, T] \times D \times D$ with $\delta_D(x) \wedge \delta_D(y) \geq a\Phi^{-1}(t) \geq 4|x - y|$.

Proof. We fix $(t, x, y) \in (0, T] \times D \times D$ satisfying $\delta_D(x) \wedge \delta_D(y) \geq a\Phi^{-1}(t) \geq 4|x - y|$. Note that $|x - y| \leq a\Phi^{-1}(t)/4 \leq a\Phi^{-1}(T)/4$ and that

$$B(x, a\Phi^{-1}(t)/4) \subset B(y, a\Phi^{-1}(t)/2) \subset B(y, 2a\Phi^{-1}(t)/3) \subset D.$$

So by the parabolic Harnack inequality [9, Theorem 5.2], there exists $c_1 = c_1(T, \gamma, \phi) > 0$ such that

$$c_1\, p_D(t/2, x, w) \;\leq\; p_D(t, x, y) \quad \text{for every } w \in B(x, a\Phi^{-1}(t)/4).$$

This together with Lemma 3.2 and (2.4) yields that

$$\begin{aligned}
p_D(t, x, y) &\geq \frac{c_1}{|B(x, a\Phi^{-1}(t)/4)|} \int_{B(x,a\Phi^{-1}(t)/4)} p_D(t/2, x, w)dw \\
&\geq c_2(\Phi^{-1}(t))^{-d} \int_{B(x,a\Phi^{-1}(t)/4)} p_{B(x,a\Phi^{-1}(t)/4)}(t/2, x, w)dw \\
&= c_2(\Phi^{-1}(t))^{-d}\, \mathbb{P}_x\left(\tau_{B(x,a\Phi^{-1}(t)/4)} > t/2\right) \;\geq\; c_3\,(\Phi^{-1}(t))^{-d},
\end{aligned}$$

where $c_i = c_i(T, a, \gamma, \phi) > 0$ for $i = 2, 3$. □

Lemma 3.4 *Let $T > 0$ and $a > 0$ be constants. There exists $c = c(a, T, \gamma, \phi) > 0$ so that*

$$\mathbb{P}_x\left(X_t^D \in B\big(y, a\Phi^{-1}(t)/2\big)\right) \geq c\,(\Phi^{-1}(t))^d\, tJ(x,y)$$

for every $(t, x, y) \in (0, T] \times D \times D$ with $\delta_D(x) \wedge \delta_D(y) \geq a\Phi^{-1}(t)$ and $a\Phi^{-1}(t) \leq 4|x - y|$.

Proof. It follows from Lemma 3.2 that, starting at $z \in B(y, a\Phi^{-1}(t)/4)$, with probability at least $c_1 = c_1(a, T, \gamma, \phi) > 0$ the process X does not move more than $a\Phi^{-1}(t)/6$ by time t. Thus, it suffices to show that there exists $c_2 = c_2(a, T, \gamma, \phi) > 0$ such that for $(t, x, y) \in (0, T] \times D \times D$ with $\delta_D(x) \wedge \delta_D(y) \geq a\Phi^{-1}(t)$ and $a\Phi^{-1}(t) \leq 4|x - y|$,

$$\mathbb{P}_x\left(X^D \text{ hits the ball } B(y, a\Phi^{-1}(t)/4) \text{ by time } t\right) \geq c_2\,(\Phi^{-1}(t))^d\, tJ(x,y). \tag{3.4}$$

Let $B_x^t := B(x, a\Phi^{-1}(t)/9)$, $B_y^t := B(y, a\Phi^{-1}(t)/9)$ and $\tau_x^t := \tau_{B_x^t}$. It follows from Lemma 3.2 and (2.4) that there exists $c_3 = c_3(a, T, \gamma, \phi) > 0$ such that

$$\mathbb{E}_x\left[t \wedge \tau_x^t\right] \geq t\,\mathbb{P}_x\left(\tau_x^t \geq t\right) \geq c_3\, t \qquad \text{for all } t \in (0, T]. \tag{3.5}$$

Since $B_x^t \cap B_y^t = \emptyset$, by the Lévy system of X and (1.5),

$$\begin{aligned}
& \mathbb{P}_x\left(X^D \text{ hits the ball } B(y, a\Phi^{-1}(t)/4) \text{ by time } t\right) \\
\geq\ & \mathbb{P}_x\left(X_{t\wedge\tau_x^t} \in B(y, a\Phi^{-1}(t)/4)\right) \geq \mathbb{E}_x\left[\int_0^{t\wedge\tau_x^t}\int_{B_y^t} J_X(X_s, u)\,du\,ds\right] \\
\geq\ & c_4\mathbb{E}_x\left[\int_0^{t\wedge\tau_x^t}\int_{B_y^t} J(X_s, u)\,du\,ds\right].
\end{aligned} \tag{3.6}$$

We consider two cases separately.

(i) Suppose $|x - y| \leq a\Phi^{-1}(T)$. Since $|x - y| \geq a\Phi^{-1}(t)/4$, we have for $s < \tau_x^t$ and $u \in B_y^t$,

$$|X_s - u| \leq |X_s - x| + |x - y| + |y - u| \leq 2|x - y|.$$

Thus by (3.5) and (3.6),

$$\begin{aligned}
& \mathbb{P}_x\left(X^D \text{ hits the ball } B(y, a\Phi^{-1}(t))/4 \text{ by time } t\right) \\
\geq\ & c_4\mathbb{E}_x\left[t \wedge \tau_x^t\right]\, |B_y^t|\, j(2|x-y|) \geq c_5\,(\Phi^{-1}(t))^d\, tj(2|x-y|)
\end{aligned}$$

for some positive constant $c_5 = c_5(a, T, \gamma, \phi) > 0$. Therefore, in view of (2.4), the assertion of the lemma holds when $|x - y| \leq a\Phi^{-1}(T)$.

(ii) Suppose $|x - y| > a\Phi^{-1}(T)$. In this case, for $s < \tau_x^t$ and $u \in B_y^t$,

$$|X_s - u| \leq |X_s - x| + |x - y| + |y - u| \leq |x - y| + a\Phi^{-1}(t)/4 \leq |x - y| + a\Phi^{-1}(T)/4.$$

Thus from (3.6) and then (3.5),

$$\begin{aligned}
& \mathbb{P}_x\left(X^D \text{ hits the ball } B(y, a\Phi^{-1}(t)/4) \text{ by time } t\right) \\
\geq\ & c_6\mathbb{E}_x\left[t \wedge \tau_x^t\right]\int_{B_y^t} j\left(|x-y| + a\Phi^{-1}(T)/4\right) du \\
\geq\ & c_7\, t\, |B_y^t|\, j\left(|x-y| + a\Phi^{-1}(T)/4\right) \\
\geq\ & c_8\, t(\Phi^{-1}(t))^d j\left(|x-y| + a\Phi^{-1}(T)/4\right)
\end{aligned}$$

for some constants $c_i = c_i(a, T, \gamma, \phi) > 0$, $i = 6, 7, 8$. Thus we conclude from (2.3) and (2.4) that the assertion of the lemma holds for $|x - y| > a\Phi^{-1}(T)$ as well. $\Box$

Proposition 3.5 *Let T and a be positive constants. There exists $c = c(T, a, \gamma, \phi) > 0$ such that*

$$p_D(t, x, y) \ \geq \ c\, tJ(x, y)$$

for every $(t, x, y) \in (0, T] \times D \times D$ with $\delta_D(x) \wedge \delta_D(y) \geq a\Phi^{-1}(t)$ and $a\Phi^{-1}(t) \leq 4|x - y|$.

Proof. By the semigroup property, Proposition 3.3, Lemma 3.4 and (2.4), there exist positive constants $c_1 = c_1(T, a, \gamma, \phi)$ and $c_2 = c_2(T, a, \gamma, \phi)$ such that

$$\begin{aligned} p_D(t, x, y) &= \int_D p_D(t/2, x, z) p_D(t/2, z, y) dz \\ &\geq \int_{B(y, a\Phi^{-1}(t/2)/2)} p_D(t/2, x, z) p_D(t/2, z, y) dz \\ &\geq c_1 (\Phi^{-1}(t/2))^{-d}\, \mathbb{P}_x \left(X^D_{t/2} \in B(y, a\Phi^{-1}(t/2)/2) \right) \\ &\geq c_2\, tJ(x, y). \end{aligned}$$

$\Box$

Combining Propositions 3.3 and 3.5, we have the following preliminary lower bound for $p_D(t, x, y)$.

Proposition 3.6 *Let T and a be positive constants. There exists $c = c(T, a, \gamma, \phi) > 0$ such that*

$$p_D(t, x, y) \ \geq \ c\, ((\Phi^{-1}(t))^{-d} \wedge tJ(x, y))$$

for every $(t, x, y) \in (0, T] \times D \times D$ with $\delta_D(x) \wedge \delta_D(y) \geq a\Phi^{-1}(t)$.

In particular, Proposition 3.6 with $D = \mathbb{R}^d$ gives

Theorem 3.7 *For any constant $T > 0$, there exists $c = c(T, \gamma, \phi) > 0$ such that for all $(t, x, y) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$,*

$$p(t, x, y) \geq c \left((\Phi^{-1}(t))^{-d} \wedge tJ(x, y) \right).$$

4 Factorization of Dirichlet heat kernel

In this section, we continue assuming that X is a purely discontinuous rotationally symmetric Lévy process with Lévy exponent Ψ and Lévy density J_X satisfying (1.6) and (1.5) respectively, where the complete Bernstein function ϕ satisfies the growth condition **(A)**.

Throughout this section, $T > 0$ is a fixed constant and D is a fixed κ-fat open set with characteristics (R_1, κ). Recall that $A_r(x) \in D$ is defined in Definition 1.2. For $(t, x) \in (0, T] \times \overline{D}$, set $r = r(t) = \Phi^{-1}(t) R_1 / \Phi^{-1}(T) \leq R_1$ and define

$$U(x, t) := D \cap B\big(x, |x - A_r(x)| + \kappa r/3\big), \quad V(x, t) := D \cap B\big(x, |x - A_r(x)| + \kappa r\big). \tag{4.1}$$



Figure 1: $U(x,t)$ and $V(x,t)$

Let

$$A'_r(x) \in D \text{ be a point such that } B\big(A'_r(x), \kappa r/3\big) \subset B\big(A_r(x), \kappa r\big) \setminus U(x,t). \tag{4.2}$$

Note that $B(A_r(x), \kappa r/3) \subset U(x,t)$ and $B(A'_r(x), \kappa\, r/6) \subset B(A'_r(x), \kappa\, r/3) \subset V(x,t) \setminus U(x,t)$. See Figure 1.

Lemma 4.1 *For every $T > 0$ and $M \geq 1$, we have that, for $(t,x) \in (0,T] \times D$,*

$$\begin{aligned} &\mathbb{P}_x(\tau_D > t/M) \asymp \mathbb{P}_x(\tau_{V(x,t)} > Mt) \asymp \mathbb{P}_x(\tau_{V(x,t)} > t/M) \asymp \mathbb{P}_x(\tau_D > Mt) \\ \asymp\ &\mathbb{P}_x(X_{\tau_{U(x,t)}} \in D) \asymp t^{-1}\mathbb{E}_x[\tau_{U(x,t)}], \end{aligned} \tag{4.3}$$

where $U(x,t)$ and $V(x,t)$ are the sets defined in (4.1) *and the (implicit) comparison constants in* (4.3) *depend only on $d, M, T, R_1, \kappa, \gamma$ and ϕ.*

Proof. Without loss of generality we assume $R_1 \leq 1$. Fix $(t,x) \in (0,T] \times D$, and set $r = r(t) = \Phi^{-1}(t) R_1 / \Phi^{-1}(T) \leq R_1 \leq 1$. Recall that $U(x,t)$ and $V(x,t)$ are the sets defined in (4.1). Observe that

$$\mathbb{P}_x(\tau_{V(x,t)} > Mt) \,\leq\, \mathbb{P}_x(\tau_{V(x,t)} > t/M) \wedge \mathbb{P}_x(\tau_D > Mt) \,\leq\, \mathbb{P}_x(\tau_D > t/M). \tag{4.4}$$

Note that by (1.5), (2.2) and (2.10), we have

$$\begin{aligned}\mathbb{P}_x\big(X(\tau_{U(x,t)}) \in B(A'_r(x), \kappa r/6)\big) &= \mathbb{E}_x \int_0^{\tau_{U(x,t)}} \int_{B(A'_r(x),\kappa r/6)} J_X(X_t, y)dtdy \\ &\asymp r^d j(r) \mathbb{E}_x\big[\tau_{U(x,t)}\big] \asymp \Phi(r)^{-1}\, \mathbb{E}_x\big[\tau_{U(x,t)}\big] \asymp t^{-1}\, \mathbb{E}_x[\tau_{U(x,t)}]. \end{aligned} \tag{4.5}$$

If $|x - A_r(x)| < \kappa r/2$, then $B(x, \kappa r/3) \subset U(x,t) \subset V(x,t)$ and so

$$1 \,\ge\, \mathbb{P}_x(\tau_D > t/M) \,\ge\, \mathbb{P}_x(\tau_D > Mt) \,\ge\, \mathbb{P}_x(\tau_{V(x,t)} > Mt) \,\ge\, \mathbb{P}_x(\tau_{B(x,\kappa r/3)} > Mt),$$

which is greater than or equal to a positive constant depending only on $\phi, T, R_1, \gamma, d, \kappa, M$ by Lemma 3.2. Thus we have, in view of (4.4),

$$1 \asymp \mathbb{P}_x(\tau_D > t/M) \asymp \mathbb{P}_x(\tau_{V(x,t)} > Mt) \asymp \mathbb{P}_x(\tau_{V(x,t)} > t/M) \asymp \mathbb{P}_x(\tau_D > Mt).$$

Moreover, when $|x - A_r(x)| < \kappa r/2$, we have $B(x, \kappa r/3) \subset U(x,t) \subset B(x,r)$ and so by Lemma 3.1 and (2.4),

$$c_1 t \le c_2 \Phi(\kappa r/3) \le \mathbb{E}_x[\tau_{B(x,\kappa r/3)}] \le \mathbb{E}_x[\tau_{U(x,t)}] \le \mathbb{E}_x[\tau_{B(x,r)}] \le c_3 \Phi(r) \le c_4 t.$$

Combining the last two displays with (4.5) and the fact that

$$1 \ge \mathbb{P}_x(X_{\tau_{U(x,t)}} \in D) \ge \mathbb{P}_x\big(X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\big),$$

we arrive at the assertion of the lemma when $|x - A_r(x)| < \kappa r/2$.

Now we assume that $|x - A_r(x)| \ge \kappa r/2$. We note that

$$\mathbb{P}_x(\tau_D > t/M) \le \mathbb{P}_x(\tau_{U(x,t)} > t/M) + \mathbb{P}_x(X_{\tau_{U(x,t)}} \in D)\,. \tag{4.6}$$

For $r \in (0, R_1]$, by Theorem 2.1, we have

$$\begin{aligned}\mathbb{P}_x\big(X_{\tau_{U(x,t)}} \in D\big) \;&\le\; c_5 \mathbb{P}_{A_r(x)}\big(X_{\tau_{U(x,t)}} \in D\big) \frac{\mathbb{P}_x\big(X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\big)}{\mathbb{P}_{A_r(x)}\big(X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\big)} \\ &\le\; c_5 \frac{\mathbb{P}_x\big(X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\big)}{\mathbb{P}_{A_r(x)}\big(X_{\tau_{B(A_r(x),\kappa r/3)}} \in B(A'_r(x), \kappa r/6)\big)}.\end{aligned}$$

Note that when $(w, y) \in B(A_r(x), \kappa r/3) \times B(A'_r(x), \kappa r/6)$, $|w-y| \le 2\kappa r$ and so $j(|w-y|) \ge j(2\kappa r)$. Thus, by (1.5), (2.10) and Lemma 3.1,

$$\begin{aligned}&\mathbb{P}_{A_r(x)}\big(X_{\tau_{B(A_r(x),\kappa r/3)}} \in B(A'_r(x), \kappa r/6)\big) = \mathbb{E}_{A_r(x)} \int_{B(A'_r(x),\kappa r/6)} \int_0^{\tau_{B(A_r(x),\kappa r/3)}} J_X(X_t, z)dtdz \\ &\ge c_6 r^d\, j(r)\, \mathbb{E}_{A_r(x)}\big[\tau_{B(A_r(x),\kappa r/3)}\big] \ge c_7 \Phi(r)^{-1}\, \mathbb{E}_0\big[\tau_{B(0,\kappa r/3)}\big] \ge c_8 \Phi(r)^{-1}\, \Phi(\kappa r/3) \ge c_9,\end{aligned}$$

where the last inequality is due to (2.4). Therefore,

$$\mathbb{P}_x\big(X_{\tau_{U(x,t)}} \in D\big) \;\le\; c_{10}\, \mathbb{P}_x\big(X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\big). \tag{4.7}$$

Since $\mathbb{P}_x\big(\tau_{U(x,t)} > t/M\big) \le Mt^{-1}\mathbb{E}_x\big[\tau_{U(x,t)}\big]$, we have by (4.6), (4.7) and (4.5),

$$\mathbb{P}_x\big(\tau_D > t/M\big) \le c_{11} t^{-1} \mathbb{E}_x\big[\tau_{U(x,t)}\big]. \tag{4.8}$$

On the other hand, by the strong Markov property of X,

$$\begin{aligned}
& \mathbb{P}_x\big(X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\big) \\
= \;& \mathbb{E}_x\left[\mathbb{P}_{X_{\tau_{U(x,t)}}}\left(\tau_{B(X_{\tau_{U(x,t)}},\kappa r/6)} > Mt\right); X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\right] \\
& + \mathbb{E}_x\left[\mathbb{P}_{X_{\tau_{U(x,t)}}}\left(\tau_{B(X_{\tau_{U(x,t)}},\kappa r/6)} \le Mt\right) : X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\right] \\
\le \;& \mathbb{E}_x\left[\mathbb{P}_{X_{\tau_{U(x,t)}}}\left(\tau_{V(x,t)} > Mt\right); X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\right] \\
& + \mathbb{P}_0\left(\tau_{B(0,\kappa r/6)} \le Mt\right) \mathbb{P}_x\left(X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\right) \\
= \;& \mathbb{P}_x\left(\tau_{V(x,t)} > Mt, X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\right) \\
& + \left(1 - \mathbb{P}_0\left(\tau_{B(0,\kappa r/6)} > Mt\right)\right) \mathbb{P}_x\left(X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\right) \\
\le \;& \mathbb{P}_x\left(\tau_{V(x,t)} > Mt\right) + \left(1 - \mathbb{P}_0\left(\tau_{B(0,\kappa r/6)} > Mt\right)\right) \mathbb{P}_x\left(X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\right).
\end{aligned}$$

Thus

$$\mathbb{P}_0\big(\tau_{B(0,\kappa r/6)} > Mt\big)\mathbb{P}_x\left(X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)\right) \le \mathbb{P}_x\left(\tau_{V(x,t)} > Mt\right). \tag{4.9}$$

Since $\mathbb{P}_0\big(\tau_{B(0,\kappa r/6)} > Mt\big) > c_{12}$ by (2.4) and Lemma 3.2, combining (4.5) with (4.9), we get

$$t^{-1}\mathbb{E}_x[\tau_{U(x,t)}] \le c_{13}\mathbb{P}_x(X_{\tau_{U(x,t)}} \in B(A'_r(x), \kappa r/6)) \le c_{14}\mathbb{P}_x(\tau_{V(x,t)} > Mt).$$

This together with (4.8) and (4.4) completes the proof of this lemma. □

Lemma 4.2 *Suppose that U_1, U_3, E are open subsets of $\mathbb{R}^d$ with $U_1, U_3 \subset E$ and $\mathrm{dist}(U_1, U_3) > 0$. Let $U_2 := E \setminus (U_1 \cup U_3)$. If $x \in U_1$ and $y \in U_3$, then for all $t > 0$,*

$$p_E(t,x,y) \le \mathbb{P}_x\left(X_{\tau_{U_1}} \in U_2\right)\left(\sup_{s<t,\, z\in U_2} p_E(s,z,y)\right) + \left(t \wedge \mathbb{E}_x\left[\tau_{U_1}\right]\right)\left(\sup_{u\in U_1,\, z\in U_3} J_X(u,z)\right) \tag{4.10}$$

and

$$p_E(t,x,y) \ \ge\ t\,\mathbb{P}_x(\tau_{U_1} > t)\,\mathbb{P}_y(\tau_{U_3} > t) \inf_{u\in U_1,\, z\in U_3} J_X(u,z). \tag{4.11}$$

Proof. For (4.10), see the proof of [15, Lemma 3.4]. For (4.11), see the proof of [16, Lemma 3.3]. □

Proof of Theorem 1.3(ii). It follows from Lemma 4.1 and the semigroup property that it suffices to establish the assertion for $T \le 1$. So in the remainder of this proof, we assume that $T \le 1$. Fix $t \in (0, T]$ and set $r = \Phi^{-1}(t)R_1/\Phi^{-1}(T)$. By (2.2), (2.4), (2.5) and Theorem 3.7,

$$p(t/2, x, y) \asymp (\Phi^{-1}(t))^{-d} \quad \text{if } |x-y| \le 8r. \tag{4.12}$$

By the semigroup property, (2.4), (2.5) and Lemma 4.1, when $|x-y| \le 8r$,

$$\begin{aligned}
p_D(t/2,x,y) &= \int_D p_D(t/4,x,z)p_D(t/4,z,y)dz \\
&\le \sup_{z\in\mathbb{R}^d} p(t/4,z,y)\mathbb{P}_x(\tau_D > t/4) \\
&\le c_1(\Phi^{-1}(t))^{-d}\mathbb{P}_x(\tau_D > t).
\end{aligned} \tag{4.13}$$

Thus by (4.12) and (4.13),

$$p_D(t/2,x,y) \le c_2\,\mathbb{P}_x(\tau_D > t)p(t/2,x,y) \quad \text{if } |x-y| \le 8r\,. \tag{4.14}$$

Now we assume $|x-y| > 8r$. Let $D_1 := U(x,t)$ be the set defined in (4.1), $D_3 := \{z \in D : |z-x| > |x-y|/2\}$ and

$$D_2 := D \setminus (D_1 \cup D_3) = \{z \in D \setminus U(x,t) : |z-x| \le |x-y|/2\}.$$

Then by condition **(B)** we have

$$\sup_{s<t/2,\, z\in D_2} p(s,z,y) \le \sup_{s<t/2,\, |z-y|\ge|x-y|/2} p(s,|z-y|) \le C_1 \sup_{s<t/2} p(s, C_2|x-y|/2).$$

Extend the definition of $p(t,r)$ by setting $p(t,r)=0$ for $t<0$ and $r\ge 0$. For each fixed $x,y\in\mathbb{R}^d$ and $t>0$ with $|x-y|>8r$, one can easily check, in view of (1.9), that $(s,w)\mapsto p(s,C_2|w-y|/2)$ is a parabolic function in $(-\infty,T]\times B(x,2r)$. So by the parabolic Harnack inequality from [9, Theorem 5.2], there is a constant $c_3=c_3(d,\gamma,\phi)\ge 1$ so that for every $t\in(0,T]$,

$$\sup_{s<t/2} p(s,C_2|x-y|/2) \le c_3 p(t/2, C_2|x-y|/2).$$

Hence we have

$$\sup_{s<t/2,\, z\in D_2} p(s,z,y) \le c_3 C_1 p(t/2, C_2|x-y|/2) \le c_3 C_1^2 p(t/2, C_2^2|x-y|/4), \tag{4.15}$$

where in the last inequality, we applied condition **(B)** again. If $u\in D_1$, then $|x-u|\le|x-A_r(x)|+\kappa r/3$. Thus if $u\in D_1$, $z\in D_3$, then

$$|u-z| \ge |z-x|-|x-u| \ge |z-x|-|x-A_r(x)|-\kappa r/3 \ge |z-x|-r \ge \frac{1}{2}|z-x| \ge \frac{1}{4}|x-y|.$$

Thus by (2.2), (1.5), (2.4) and Theorem 3.7,

$$\begin{aligned} t\sup_{u\in D_1,\, z\in D_3} J_X(u,z) &\le \gamma t \sup_{|u-z|\ge|x-y|/4} j(|u-z|) \le \gamma t j(|x-y|/4) \le \gamma t j(C_2^2|x-y|/4) \\ &\le c_4 p(t/2, C_2^2|x-y|/4). \end{aligned}$$

Let $C_5 := C_2^2/4$. It follows then from (4.15), Lemmas 4.1 and 4.2 that for $|x-y|>8r$,

$$\begin{aligned} p_D(t/2,x,y) &\le c_5 p(t/2, C_5x, C_5y)\left(\mathbb{P}_x(X_{\tau_{U(x,t)}}\in D) + t^{-1}\mathbb{E}_x[\tau_{U(x,t)}]\right) \\ &\le c_6 \mathbb{P}_x(\tau_D>t)\, p(t/2, C_5x, C_5y). \end{aligned} \tag{4.16}$$

Hence by condition **(B)**, (4.14), (4.16), symmetry, the semigroup property and Lemma 4.1, we conclude that for all $(t,x,y)\in(0,T]\times D\times D$,

$$\begin{aligned} p_D(t,x,y) &= \int_D p_D(t/2,x,z)p_D(t/2,z,y)dz \\ &\le c_6^2 \mathbb{P}_x(\tau_D>t)\mathbb{P}_y(\tau_D>t)\int_{\mathbb{R}^d} p(t/2,C_5x,C_5z)p(t/2,C_5z,C_5y)dz \\ &= c_7 \mathbb{P}_x(\tau_D>t)\mathbb{P}_y(\tau_D>t)\int_{\mathbb{R}^d} p(t/2,C_5x,z)p(t/2,z,C_5y)dz \\ &= c_7 \mathbb{P}_x(\tau_D>t)\mathbb{P}_y(\tau_D>t)p(t,C_5x,C_5y)\,. \end{aligned}$$

□

Proof of Theorem 1.3(i). Fix $(t,x,y) \in (0,T] \times D \times D$ and set $r = \Phi^{-1}(t)R_1/\Phi^{-1}(T)$. Let $U(x,t)$ be the set defined in (4.1), $A_r(x) \in D$ and $A'_r(x) \in D$ be the points defined in Definition 1.2 and (4.2) respectively. Note that

$$\begin{aligned} p_D(t,x,y) &= \int_{D\times D} p_D(t/3,x,u)p_D(t/3,u,v)p_D(t/3,u,y)dudv \\ &\geq \inf_{(u,v)\in B(A'_r(x),\kappa r/6)\times B(A'_r(y),\kappa r/6)} p_D(t/3,u,v)\int_{B(A'_r(x),\kappa r/6)} p_D(t/3,x,u)du \\ &\quad \cdot \int_{B(A'_r(y),\kappa r/6)} p_D(t/3,v,y)dv. \end{aligned} \tag{4.17}$$

For $(u,v) \in B(A'_r(x),\kappa r/6) \times B(A'_r(y),\kappa r/6)$,

$$\begin{aligned} &\inf_{(u,v)\in B(A'_r(x),\kappa r/6)\times B(A'_r(y),\kappa r/6)} p_D(t/3,u,v) \\ \geq\ & c_1 \inf_{(u,v)\in B(A'_r(x),\kappa r/6)\times B(A'_r(y),\kappa r/6)} \Big((\Phi^{-1}(t/3))^{-d} \wedge (t/3)J(u,v)\Big) \\ \geq\ & c_2(tJ(x,y) \wedge (\Phi^{-1}(t))^{-d}), \end{aligned} \tag{4.18}$$

where in the first inequality we used Proposition 3.6, in the second inequality we used (2.2) and (2.3) respectively in the cases $|x-y| \geq \kappa r$ and $|x-y| < \kappa r$.

On the other hand, for $u \in B(A'_r(x),\kappa r/6)$, by Lemma 4.2 with $U_1 = U(x,t)$ and $U_3 = B(A'_r(x),\kappa r/4)$, we have

$$\begin{aligned} p_D(t/3,x,u) &\geq t\mathbb{P}_x(\tau_{U(x,t)} > t/3)\mathbb{P}_u(\tau_{U_3} > t/3) \inf_{w\in U(x,t),\, z\in U_3} J_X(w,z) \\ &\geq \gamma^{-1}t\mathbb{P}_x(\tau_{U(x,t)} > t/3)\mathbb{P}_u(\tau_{B(u,\kappa r/16)} > t/3) \inf_{w\in U(x,t),\, z\in U_3} j(|w-z|). \end{aligned}$$

Since

$$j(|w-z|) \geq j(r) = j(\Phi^{-1}(t)R_1/\Phi^{-1}(T)) \quad \text{for } (w,z) \in U(x,t)\times U_3,$$

we have by Lemma 3.2, (2.2) and (2.4),

$$p_D(t/3,x,u) \geq c_3 t\,\mathbb{P}_x(\tau_{U(x,t)} > t/3)\frac{1}{\Phi^{-1}(t)^d\, t} = c_3\Phi^{-1}(t)^{-d}\,\mathbb{P}_x(\tau_{U(x,t)} > t/3).$$

It follows from condition **(A)** that there exists $c_4 > 1$ such that $\Phi^{-1}(bt) \leq c_4 b^{1/(2\delta_2)}\Phi^{-1}(t)$ for every $t \leq T$ and $b \in (0,1]$. Thus with $a := (\kappa/(3c_4))^{2\delta_2}$ we have that $3^{-1}\kappa\Phi^{-1}(t) \geq \Phi^{-1}(at)$ for every $t \leq T$, and hence $V(x,at) \subset U(x,t)$. Thus Lemma 4.1 implies that

$$\mathbb{P}_x(\tau_{U(x,t)} > t/3) \geq \mathbb{P}_x(\tau_{V(x,at)} > t/3) \geq c_5\mathbb{P}_x(\tau_D > t).$$

Consequently, by (2.4),

$$\int_{B(A'_r(x),\kappa r/6)} p_D(t/3,x,u)du \geq \frac{c_6}{\Phi^{-1}(t)^d}\mathbb{P}_x(\tau_D > t)|B(A'_r(x),\kappa r/6)| \geq c_7\mathbb{P}_x(\tau_D > t).$$

Similarly, using the symmetry we also have $\int_{B(A'_r(y),\kappa r/6)} p_D(t/3,v,y)dv \geq c_8\mathbb{P}_y(\tau_D > t)$. Hence we have by (4.17) and (4.18),

$$p_D(t,x,y) \geq c_9\,\mathbb{P}_x(\tau_D > t)\mathbb{P}_y(\tau_D > t)\Big((\Phi^{-1}(t))^{-d} \wedge tJ(x,y)\Big).$$

□

5 Small time Dirichlet heat kernel estimates for subordinate Brownian motions in $C^{1,1}$ open set

In this section we assume that D is a $C^{1,1}$ open set in $\mathbb{R}^d$ with characteristics (R_2, Λ) and that X is a subordinate Brownian motion with Lévy exponent $\Psi(\xi) = \phi(|\xi|^2)$, where ϕ is a complete Bernstein function satisfying condition **(A)**.

Proof of Theorem 1.5(ii). Without loss of generality we assume $R_2 \le 1$. In view of Theorem 1.3(ii), it suffices to show that

$$\mathbb{P}_x\left(\tau_D > t\right) \le c\left(1 \wedge \frac{\Phi(\delta_D(x))}{t}\right)^{1/2} \quad \text{for } (t,x) \in (0,T] \times D. \tag{5.1}$$

Fix $(t,x) \in (0,T] \times D$ and set $r = r(t) = \Phi^{-1}(t)R_2/\Phi^{-1}(T) \le R_2 \le 1$. By Theorem 3.7, we only need to show the theorem for $\delta_D(x) < r/16$. Take $x_0 \in \partial D$ such that $\delta_D(x) = |x - x_0|$.

Let $U_1 := B(x_0, r/8) \cap D$ and $\mathbf{n}(x_0)$ be the unit inward normal of ∂D at the point x_0. Put

$$x_1 = x_0 + \frac{r}{16}\mathbf{n}(x_0).$$

Note that $\delta_D(x_1) = r/16$. Applying the boundary Harnack principle (Theorem 2.3) and (2.4) we get

$$\begin{aligned}\mathbb{P}_x(X_{\tau_{U_1}} \in D \setminus U_1) &\le c_1 \mathbb{P}_{x_1}(X_{\tau_{U_1}} \in D \setminus U_1)\sqrt{\frac{\Phi(\delta_D(x))}{\Phi(\delta_D(x_1))}} \\ &\le c_2 \mathbb{P}_{x_1}(X_{\tau_{U_1}} \in D \setminus U_1)\sqrt{\frac{\Phi(\delta_D(x))}{t}} \le c_2\sqrt{\frac{\Phi(\delta_D(x))}{t}}.\end{aligned} \tag{5.2}$$

Take $x_2 \in \mathbb{R}^d$ so that $B(x_2, r) \subset B(x_0, 4r) \setminus B(x_0, r)$. Then, by (2.10), (2.2) and (2.4), we have

$$\begin{aligned}\mathbb{P}_x(X_{\tau_{U_1}} \in B(x_2, r)) &= \mathbb{E}_x\left[\int_0^{\tau_{U_1}} \int_{B(x_2,r)} J(|X_s - y|)dyds\right] \\ &\ge c_3|B(0, \Phi^{-1}(t))|j(5r)\mathbb{E}_x[\tau_{U_1}] \\ &\ge c_4\Phi(\Phi^{-1}(t))^{-1}\mathbb{E}_x[\tau_{U_1}] = c_4 t^{-1}\mathbb{E}_x[\tau_{U_1}].\end{aligned}$$

Now by the same argument as that in (5.2), we get,

$$\mathbb{E}_x[\tau_{U_1}] \le c_4^{-1} t\mathbb{P}_x(X_{\tau_{U_1}} \in B(x_2, r)) \le c_5 t\mathbb{P}_{x_1}(X_{\tau_{U_1}} \in B(x_2, r))\sqrt{\frac{\Phi(\delta_D(x))}{t}} \le c_5\sqrt{t\,\Phi(\delta_D(x))}. \tag{5.3}$$

Thus, by (5.2) and (5.3), we have

$$\begin{aligned}\mathbb{P}_x\left(\tau_D > t\right) &\le \mathbb{P}_x\left(\tau_{U_1} > t\right) + \mathbb{P}_x\left(X_{\tau_{U_1}} \in D \setminus U_1\right) \\ &\le \frac{1}{t}\mathbb{E}_x\left[\tau_{U_1}\right] + \mathbb{P}_x\left(X_{\tau_{U_1}} \in D \setminus U_1\right) \le c_6\sqrt{\frac{\Phi(\delta_D(x))}{t}}.\end{aligned} \tag{5.4}$$

Since $\mathbb{P}_x\left(\tau_D > t\right) \le 1$, (5.1) follows immediately. $\square$

Let $\delta_{\partial D}(x)$ be the Euclidean distance between x and ∂D. It is well-known that any $C^{1,1}$ open set D with $C^{1,1}$-characteristics (R_2, Λ) satisfies both the *uniform interior ball condition* and the *uniform exterior ball condition*: there exists $r_0 = r_0(R_2, \Lambda) \le R_2$ such that for every $x \in D$ with $\delta_{\partial D}(x) < r_0$ and $y \in \mathbb{R}^d \setminus \overline{D}$ with $\delta_{\partial D}(y) < r_0$, there are $z_x, z_y \in \partial D$ so that $|x - z_x| = \delta_{\partial D}(x)$, $|y - z_y| = \delta_{\partial D}(y)$ and that $B(x_0, r_0) \subset D$ and $B(y_0, r_0) \subset \mathbb{R}^d \setminus \overline{D}$ for $x_0 = z_x + r_0(x - z_x)/|x - z_x|$ and $y_0 = z_y + r_0(y - z_y)/|y - z_y|$.

In the remainder of this section, we fix such an r_0 and set $T_0 := \Phi(r_0/16)$. For any $x \in D$ with $\delta_D(x) < r_0$, let z_x be a point on ∂D such that $|z_x - x| = \delta_D(x)$ and $\mathbf{n}(z_x) := (x - z_x)/|z_x - x|$.

Lemma 5.1 *Let $\kappa_0 \in (0,1)$ and $a > 0$. There exists a constant $c = c(\kappa_0, R_2, r_0, a, \phi) > 0$ such that for every $(t, x) \in (0, T_0] \times D$ with $\delta_D(x) \le 3\Phi^{-1}(t) < r_0/4$ and $\kappa_0 \in (0,1)$,*

$$\mathbb{P}_x \left(X^D_{at} \in B(x_0, \kappa_0 \Phi^{-1}(t)) \right) \ \ge\ c \sqrt{\frac{\Phi(\delta_D(x))}{t}}, \tag{5.5}$$

where $x_0 := z_x + \frac{9}{2}\Phi^{-1}(t)\mathbf{n}(z_x)$.

Proof. Let $0 < \kappa_1 \le \kappa_0$ and assume first that $2^{-4}\kappa_1\Phi^{-1}(t) < \delta_D(x) \le 3\Phi^{-1}(t)$. As in the proof of Lemma 3.4, we get that, in this case, using the fact that $|x - x_0| \in [\frac{3}{2}\kappa_0\Phi^{-1}(t), 6\Phi^{-1}(t)]$, there exist constants $c_i = c_i(\kappa_1, r_0, a) > 0$, $i = 1, 2$, such that for all $t \le T_0$, we have

$$\mathbb{P}_x \left(X^D_{at} \in B(x_0, \kappa_1 \Phi^{-1}(t)) \right) \ \ge\ c_1 t (\Phi^{-1}(t))^d J(x, x_0) \ \ge\ c_2 > 0. \tag{5.6}$$

By taking $\kappa_1 = \kappa_0$, this shows that (5.5) holds for all $a > 0$ in the case when $2^{-4}\kappa_0\Phi^{-1}(t) < \delta_D(x) \le 3\Phi^{-1}(t)$.

So it suffices to consider the case that $\delta_D(x) \le 2^{-4}\kappa_0\Phi^{-1}(t)$. We now show that there is some $a_0 > 1$ so that (5.5) holds for every $a \ge a_0$ and $\delta_D(x) \le 2^{-4}\kappa_0\Phi^{-1}(t)$. For simplicity, we assume without loss of generality that $x_0 = 0$ and let $\widehat{B} := B(0, \kappa_0\Phi^{-1}(t))$. Let $U := D \cap B(z_x, \kappa_0\Phi^{-1}(t))$. By the strong Markov property of X^D at the first exit time τ_U from U and Lemma 3.2, there exists $c_3 = c_3(a) > 0$ such that

$$\begin{aligned}
&\mathbb{P}_x \left(X^D_{at} \in \widehat{B} \right) \\
\ge\ & \mathbb{P}_x \left(\tau_U < at,\ X_{\tau_U} \in B(0, 2^{-1}\kappa_0\Phi^{-1}(t)) \text{ and } |X^D_s - X_{\tau_U}| < 2^{-1}\kappa_0\Phi^{-1}(t) \text{ for } s \in [\tau_U, \tau_U + at] \right) \\
\ge\ & c_3\, \mathbb{P}_x \left(\tau_U < at \text{ and } X_{\tau_U} \in B(0, 2^{-1}\kappa_0\Phi^{-1}(t)) \right).
\end{aligned} \tag{5.7}$$

Let $x_1 = z_x + 4^{-1}\kappa_0\mathbf{n}(z_x)\Phi^{-1}(t)$ and $B_1 := B(x_1, 4^{-1}\kappa_0\Phi^{-1}(t))$. It follows from the boundary Harnack principle (Theorem 2.3) and (2.4) that there exist $c_k = c_k(R_2, \Lambda, \gamma, \phi) > 0$, $k = 4, 5$, such that for all $t \in (0, T_0]$,

$$\begin{aligned}
\mathbb{P}_x \left(X_{\tau_U} \in B(0, 2^{-1}\kappa_0\Phi^{-1}(t)) \right) \ &\ge\ c_4 \mathbb{P}_{x_1} \left(X_{\tau_U} \in B(0, 2^{-1}\kappa_0\Phi^{-1}(t)) \right) \sqrt{\frac{\Phi(\delta_D(x))}{\Phi(\delta_D(x_1))}} \\
&\ge\ c_5 \mathbb{P}_{x_1} \left(X_{\tau_{B_1}} \in B(0, 2^{-1}\kappa_0\Phi^{-1}(t)) \right) \sqrt{\frac{\Phi(\delta_D(x))}{t}}.
\end{aligned}$$

By (2.10), Lemma 3.1 and (2.2)-(2.4), we have

$$\begin{aligned}
&\mathbb{P}_{x_1}\left(X_{\tau_{B_1}} \in B(0,\kappa_0\Phi^{-1}(t)/2)\right) = \mathbb{E}_x\left[\int_0^{\tau_{B_1}}\int_{B(0,\kappa_0\Phi^{-1}(t)/2)} J(X_s,y)dy\right]\\
\geq\;& c_6 j(\kappa_0\Phi^{-1}(t))\,|B(0,\kappa_0\Phi^{-1}(t)/2)|\,\mathbb{E}_x\left[\tau_{B_1}\right]\\
\geq\;& \frac{c_7}{(\kappa_0\Phi^{-1}(t))^d\Phi(\kappa_0\Phi^{-1}(t))}\left(\kappa_0\Phi^{-1}(t)\right)^d\Phi(\kappa_0\Phi^{-1}(t)) = c_7.
\end{aligned}$$

Thus

$$\mathbb{P}_x\left(X_{\tau_U} \in B(0,2^{-1}\kappa_0\Phi^{-1}(t))\right) \geq c_8\sqrt{\frac{\Phi(\delta_D(x))}{t}}. \tag{5.8}$$

It follows from (5.3) that there exists $c_9 > 0$ such that

$$\mathbb{P}_x(\tau_U \geq at) \;\leq\; (at)^{-1}\,\mathbb{E}_x[\tau_U] \;\leq\; a^{-1}c_9\sqrt{\frac{\Phi(\delta_D(x))}{t}}.$$

Define $a_0 = 2c_9/(c_8)$. We have by (5.7)–(5.8) and the display above that for $a \geq a_0$,

$$\begin{aligned}
\mathbb{P}_x(X^D_{at} \in \widehat{B}) \;&\geq\; c_3\left(\mathbb{P}_x(X_{\tau_U} \in B(0,2^{-1}\kappa_0\Phi^{-1}(t))) - \mathbb{P}_x\left(\tau_U \geq at\right)\right)\\
&\geq\; c_3\,(c_9/2)\sqrt{\frac{\Phi(\delta_D(x))}{t}}.
\end{aligned} \tag{5.9}$$

(5.6) and (5.9) show that (5.5) holds for every $a \geq a_0$ and for every $x \in D$ with $\delta_D(x) \leq 3\Phi^{-1}(t)$.

Now we deal with the case $0 < a < a_0$ and $\delta_D(x) \leq 2^{-4}\kappa_0\Phi^{-1}(t)$. If $\delta_D(x) \leq 3\Phi^{-1}(at/a_0)$, we have from (5.5) for the case of $a = a_0$ that there exist $c_{10} = c_{10}(\kappa_0, R_2, \Lambda, a, \phi) > 0$ and $c_{11} = c_{11}(\kappa_0, R_2, \Lambda, a, \phi) > 0$ such that

$$\begin{aligned}
\mathbb{P}_x\left(X^D_{at} \in B(x_0,\,\kappa_0\Phi^{-1}(t))\right) \;&\geq\; \mathbb{P}_x\left(X^D_{a_0(at/a_0)} \in B(x_0,\,\kappa_0\Phi^{-1}(at/a_0))\right)\\
&\geq\; c_{10}\sqrt{\frac{\Phi(\delta_D(x))}{at/a_0}} = c_{11}\sqrt{\frac{\Phi(\delta_D(x))}{t}}.
\end{aligned}$$

If $3\Phi^{-1}(at/a_0) < \delta_D(x) \leq 2^{-4}\kappa_0\Phi^{-1}(t)$ (in this case $1 > \kappa_0 > 3\cdot 2^4\Phi^{-1}(a/a_0)$), we get (5.5) from (5.6) by taking $\kappa_1 = \Phi^{-1}(a/a_0)$. The proof of the lemma is now complete. □

Proof of Theorem 1.5(i). By Theorem 1.3(i), it suffices to show that

$$\mathbb{P}_x\left(\tau_D > t\right) \geq c\left(1 \wedge \frac{\Phi(\delta_D(x))}{t}\right)^{1/2} \quad \text{for } (t,x) \in (0,T] \times D. \tag{5.10}$$

Assume $(t,x) \in (0,T] \times D$. Since D satisfies the uniform interior ball condition with radius r_0 and $0 < (T_0/T)t \leq T_0$, we can choose a point ξ^t_x as follows: if $\delta_D(x) \leq 3\Phi^{-1}((T_0/T)t)$, let $\xi^t_x = z_x + (9/2)\Phi^{-1}((T_0/T)t)\mathbf{n}(z_x)$ so that

$$B(\xi^t_x, (3/2)\Phi^{-1}((T_0/T)t)) \subset B(z_x + 3\Phi^{-1}((T_0/T)t)\mathbf{n}(z_x), 3\Phi^{-1}((T_0/T)t)) \setminus \{x\}$$

and $\delta_D(z) \geq 3\Phi^{-1}((T_0/T)t)$ for every $z \in B(\xi^t_x, (3/2)\Phi^{-1}((T_0/T)t)))$. If $\delta_D(x) > 3\Phi^{-1}((T_0/T)t)$, choose $\xi^t_x \in B(x, \delta_D(x))$ so that $|x - \xi^t_x| = (3/2)\Phi^{-1}((T_0/T)t)$. Note that in this case,

$$B(\xi^t_x, (3/2)\Phi^{-1}((T_0/T)t)) \subset B(x, \delta_D(x)) \setminus \{x\}$$

and $\delta_D(z) \geq \Phi^{-1}((T_0/T)t)$ for every $z \in B(\xi_x^t, 2^{-1}\Phi^{-1}((T_0/T)t))$. We also define ξ_y^t the same way.

If $\delta_D(x) \leq 3\Phi^{-1}((T_0/T)t)$, by Lemma 4.1 (with $M = T/T_0$ when $T \geq T_0$) and Lemma 5.1 (with $a = 1, \kappa = 2^{-1}$),

$$\mathbb{P}_x(\tau_D > t) \geq c_1\mathbb{P}_x(\tau_D > (T_0/T)t) \geq c_1\mathbb{P}_x\left(X^D_{(T_0/T)t} \in B(\xi_x^t, 2^{-1}\Phi^{-1}((T_0/T)t))\right) \geq c_2\sqrt{\frac{\Phi(\delta_D(x))}{t}}.$$

If $\delta_D(x) > 3\Phi^{-1}((T_0/T)t)$, by Lemma 4.1, Proposition 3.6 and (2.4),

$$\begin{aligned} &\mathbb{P}_x(\tau_D > t) \geq c_1\mathbb{P}_x\left(X^D_{(T_0/T)t} \in B(\xi_x^t, 2^{-1}\Phi^{-1}((T_0/T)t))\right) \\ = \;& c_1\int_{B(\xi_x^t, 2^{-1}\Phi^{-1}((T_0/T)t))} p_D((T_0/T)t, x, u)du \;\geq\; c_3 \;\geq\; c_4\left(1 \wedge \frac{\Phi(\delta_D(x))}{t}\right)^{1/2}. \end{aligned}$$

□

6 Large time heat kernel estimates

In this section, we first give the proofs of Theorems 1.3(iii) and 1.5(iii).

Proof of Theorem 1.3(iii). Since D is bounded, in view of (2.5), the transition semigroup $\{P_t^D, t > 0\}$ of X^D consists of Hilbert-Schmidt operators, and hence compact operators, in $L^2(D; dx)$. So P_t^D has discrete spectrum $\{e^{-\lambda_k t}; k \geq 1\}$, arranged in decreasing order and repeated according to their multiplicity. Let $\{\phi_k, k \geq 1\}$ be the corresponding eigenfunctions with unit L^2-norm ($\|\phi_1\|_{L^2(D)} = 1$) which forms an orthonormal basis for $L^2(D; dx)$.

Clearly, for every $k \geq 1$

$$\int_D \mathbb{P}_x(\tau_D > 1)\phi_k(x)dx \leq |D|^{1/2}\|\phi_k\|_{L^2(D)} = |D|^{1/2}. \tag{6.1}$$

By using the eigenfunction expansion of p_D we get

$$\int_{D\times D} \mathbb{P}_x(\tau_D > 1)p_D(t, x, y)\mathbb{P}_y(\tau_D > 1)\,dxdy = \sum_{k=1}^{\infty} e^{-t\lambda_k}\left(\int_D \mathbb{P}_x(\tau_D > 1)\phi_k(x)dx\right)^2. \tag{6.2}$$

Noting that λ_k is increasing and $\|f\|^2_{L^2(D)} = \sum_{k=1}^{\infty}(\int_D f(z)\phi_k(z)dz)^2$, we have for all $t > 0$,

$$\begin{aligned} \int_{D\times D} \mathbb{P}_x(\tau_D > 1)p_D(t, x, y)\mathbb{P}_y(\tau_D > 1)\,dxdy &\leq e^{-t\lambda_1}\int_D \mathbb{P}_x(\tau_D > 1)^2 dx \\ &\leq e^{-t\lambda_1}|D|. \end{aligned} \tag{6.3}$$

On the other hand, by Theorem 1.3(ii), Remark 1.1(iii) and (6.1) we have that there is a constant $c_1 > 0$ so that for every $x \in D$,

$$\begin{aligned} &\phi_1(x) = e^{\lambda_1}\int_D p_D(1, x, y)\phi_1(y)dy \\ &\leq c_1\mathbb{P}_x(\tau_D > 1)\int_D \mathbb{P}_y(\tau_D > 1)\phi_1(y)dy \leq c_1|D|^{1/2}\,\mathbb{P}_x(\tau_D > 1). \end{aligned} \tag{6.4}$$

It now follows from (6.2) that for every that for every $t > 0$,

$$\int_{D\times D} \mathbb{P}_x(\tau_D > 1)p_D(t,x,y)\mathbb{P}_y(\tau_D > 1)\,dxdy \geq e^{-t\lambda_1}\left(\int_D \mathbb{P}_x(\tau_D > 1)\phi_1(x)dx\right)^2$$

$$\geq e^{-t\lambda_1}\left(\int_D c_1^{-1}|D|^{-1/2}\phi_1(x)^2dx\right)^2 = c_1^{-2}|D|^{-1}\,e^{-t\lambda_1}. \tag{6.5}$$

For $t \geq 3$ and $x, y \in D$, we have that

$$p_D(t,x,y) = \int_{D\times D} p_D(1,x,z)p_D(t-2,z,w)p_D(1,w,y)dzdw. \tag{6.6}$$

By Theorem 1.3(ii), Remark 1.1(iii), (2.5) and (6.3) we have that there are constants $c_i > 0$, $i = 2, 3$, so that for every $t \geq 3$ and $x, y \in D$,

$$p_D(t,x,y) \leq c_2\mathbb{P}_x(\tau_D > 1)\mathbb{P}_y(\tau_D > 1)\int_{D\times D} \mathbb{P}_z(\tau_D > 1)p_D(t-2,z,w)\mathbb{P}_w(\tau_D > 1)dzdw$$

$$\leq c_3\,\mathbb{P}_x(\tau_D > 1)\mathbb{P}_y(\tau_D > 1)e^{-t\lambda_1}. \tag{6.7}$$

By Theorem 1.3(i), Theorem 3.7, the boundedness of D and (6.5) we have that there are constants $c_i > 0$, $i = 4, 5$, so that for every $t \geq 3$ and $x, y \in D$,

$$p_D(t,x,y) \geq c_4\,\mathbb{P}_x(\tau_D > 1)\mathbb{P}_y(\tau_D > 1)\int_{D\times D} \mathbb{P}_z(\tau_D > 1)p_D(t-2,z,w)\mathbb{P}_w(\tau_D > 1)dzdw$$

$$\geq c_5\,\mathbb{P}_x(\tau_D > 1)\mathbb{P}_y(\tau_D > 1)e^{-t\lambda_1}.$$

This combined with (6.7) establishes Theorem 1.3(iii). □

Proof of Theorem 1.5(iii). By Theorem 1.5(i), it suffices to prove the theorem for $T \geq 3$. By (5.1), (5.10) and the boundedness of D, $\mathbb{P}_x(\tau_D > 1) \asymp \Phi(\delta_D(x))^{1/2}$. This and Theorem 1.3(iii) imply Theorem 1.5(iii). □

7 Green function estimates

In this section, we use Theorem 1.5 to get sharp two-sided estimates on the Green functions of subordinate Brownian motions in bounded $C^{1,1}$ open sets. We first establish the following two lemmas.

Lemma 7.1 *For every $r \in (0, 1]$ and every open subset U of $\mathbb{R}^d$,*

$$\frac{1}{2}\left(1 \wedge \frac{r^2\Phi(\delta_U(x))^{1/2}\Phi(\delta_U(y))^{1/2}}{\Phi(|x-y|)}\right) \leq \left(1 \wedge \frac{r\Phi(\delta_U(x))^{1/2}}{\Phi(|x-y|)^{1/2}}\right)\left(1 \wedge \frac{r\Phi(\delta_U(y))^{1/2}}{\Phi(|x-y|)^{1/2}}\right)$$

$$\leq 1 \wedge \frac{r^2\Phi(\delta_U(x))^{1/2}\Phi(\delta_U(y))^{1/2}}{\Phi(|x-y|)}. \tag{7.1}$$

Proof. The second inequality holds trivially. Without loss of generality, we assume $\delta_U(x) \le \delta_U(y)$. If both $\frac{r\Phi(\delta_U(x))^{1/2}}{\Phi(|x-y|)^{1/2}}$ and $\frac{r\Phi(\delta_U(y))^{1/2}}{\Phi(|x-y|)^{1/2}}$ are less than 1 or if both are larger than one,

$$\left(1 \wedge \frac{r\Phi(\delta_U(x))^{1/2}}{\Phi(|x-y|)^{1/2}}\right)\left(1 \wedge \frac{r\Phi(\delta_U(y))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) = 1 \wedge \frac{r^2\Phi(\delta_U(x))^{1/2}\Phi(\delta_U(y))^{1/2}}{\Phi(|x-y|)}.$$

So we only need to consider the case when $\frac{r\Phi(\delta_U(x))^{1/2}}{\Phi(|x-y|)^{1/2}} \le 1 < \frac{r\Phi(\delta_U(y))^{1/2}}{\Phi(|x-y|)^{1/2}}$. Note that $\Phi(\delta_U(y)) \le \Phi(\delta_U(x) + |x-y|)$. If $\delta_U(x) \ge |x-y|$, then by (1.8), $\Phi(\delta_U(y)) \le \Phi(2\delta_U(x)) \le 4\Phi(\delta_U(x))$ and so

$$1 \wedge \frac{r^2\Phi(\delta_U(x))^{1/2}\Phi(\delta_U(y))^{1/2}}{\Phi(|x-y|)} \le 1 \wedge \frac{2r\Phi(\delta_U(x))}{\Phi(|x-y|)} \le 2\left(1 \wedge \frac{r\Phi(\delta_U(x))^{1/2}}{\Phi(|x-y|)^{1/2}}\right).$$

When $\delta_U(x) < |x-y|$, by (1.8) again, $\Phi(\delta_U(y)) \le \Phi(2|x-y|) \le 4\Phi(|x-y|)$ and so

$$1 \wedge \frac{r^2\Phi(\delta_U(x))^{1/2}\Phi(\delta_U(y))^{1/2}}{\Phi(|x-y|)} \le 1 \wedge \frac{2r^2\Phi(\delta_U(x))^{1/2}}{\Phi(|x-y|)^{1/2}} \le 2\left(1 \wedge \frac{r\Phi(\delta_U(x))^{1/2}}{\Phi(|x-y|)^{1/2}}\right),$$

where the assumption $r \le 1$ is used in the last inequality. This establishes the first inequality of (7.1). □

By condition **(A)**, we have that for every $T > 0$, there exist $C_T > 1$ such that

$$C_T^{-1}\left(\frac{r}{R}\right)^{1/(2\delta_1)} \le \frac{\Phi^{-1}(r)}{\Phi^{-1}(R)} \le C_T\left(\frac{r}{R}\right)^{1/(2\delta_2)} \quad \text{for } 0 < r \le R \le T. \tag{7.2}$$

Moreover, for every $M > 0$, we have

$$r\Phi'(r) \asymp \Phi(r) \quad \text{for } r \in (0, M] \tag{7.3}$$

(see the paragraph after [19, Lemma 1.3]).

Lemma 7.2 *Suppose $T > 0$ and set*

$$h_T(a,r) = a + \Phi(r)\int_{\Phi(r)/T}^{1}\left(1 \wedge \frac{ua}{\Phi(r)}\right)\frac{1}{u^2\Phi^{-1}(u^{-1}\Phi(r))}\,du + \frac{\Phi(r)}{r}\left(1 \wedge \frac{a}{\Phi(r)}\right). \tag{7.4}$$

Then

$$h_T(a,r) \asymp \frac{a}{r} \wedge \left(\frac{a}{\Phi^{-1}(a)} + \left(\int_r^{\Phi^{-1}(a)}\frac{\Phi(s)}{s^2}ds\right)^+\right)$$

for $0 < r \le \Phi^{-1}(T/2)$ and $0 < a \le (2^{-1} \wedge (2C_T)^{-2\delta_2})T$, where C_T is the constant in (7.2) and $x^+ := x \vee 0$.

Proof. For (a,r) with $0 < a < \Phi(r) \le T/2$,

$$h_T(a,r) \asymp a + a\int_{\Phi(r)/T}^{1}\frac{du}{u\Phi^{-1}(u^{-1}\Phi(r))} + \frac{a}{r} = a + \frac{a}{r}\int_{\Phi(r)/T}^{1}\frac{\Phi^{-1}(\Phi(r))}{\Phi^{-1}(u^{-1}\Phi(r))}u^{-1}du + \frac{a}{r}.$$

By (7.2), since $\Phi(r) \le T/2$, we have

$$0 < c_2 = c_1^{-1} \int_{1/2}^{1} u^{\frac{1}{2\delta_1}-1} du \le \int_{\Phi(r)/T}^{1} \frac{\Phi^{-1}(\Phi(r))}{\Phi^{-1}(u^{-1}\Phi(r))} u^{-1} du \le c_1 \int_0^1 u^{\frac{1}{2\delta_2}-1} du = c_3 < \infty.$$

Thus, for $0 < a < \Phi(r) \le T/2$, we have

$$c_2 \frac{a}{r} \le h_T(a,r) \le c_3 \left(a + \frac{a}{r}\right) \le c_4 \frac{a}{r}. \tag{7.5}$$

On the other hand, for (a,r) with $\Phi(r) \le a \le (2^{-1} \wedge (2C_T)^{-2\delta_2})T$, using the change of variable $u = \Phi(r)/\Phi(s)$ and then applying integration by parts for the first integral below, we have

$$\begin{aligned} h_T(a,r) &\asymp a + \Phi(r) \int_{\Phi(r)/a}^{1} \frac{du}{u^2 \Phi^{-1}(u^{-1}\Phi(r))} + a \int_{\Phi(r)/T}^{\Phi(r)/a} \frac{du}{u \Phi^{-1}(u^{-1}\Phi(r))} + \frac{\Phi(r)}{r} \\ &= a + \int_r^{\Phi^{-1}(a)} \frac{\Phi'(s)}{s} ds + a \int_{\Phi^{-1}(a)}^{\Phi^{-1}(T)} \frac{\Phi'(s)}{s\Phi(s)} ds + \frac{\Phi(r)}{r} \\ &= a + \left(\frac{a}{\Phi^{-1}(a)} - \frac{\Phi(r)}{r} \right) + \int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2} ds + a \int_{\Phi^{-1}(a)}^{\Phi^{-1}(T)} \frac{\Phi'(s)}{s\Phi(s)} ds + \frac{\Phi(r)}{r} \\ &= a + \frac{a}{\Phi^{-1}(a)} + \int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2} ds + a \int_{\Phi^{-1}(a)}^{\Phi^{-1}(T)} \frac{\Phi'(s)}{s\Phi(s)} ds. \end{aligned} \tag{7.6}$$

Since $a \le \left(2^{-1} \wedge (2C_T)^{-2\delta_2}\right) T$, by (7.2) and the fact that Φ^{-1} is increasing,

$$\frac{1}{\Phi^{-1}(a)} - \frac{1}{\Phi^{-1}(T)} \asymp \frac{1}{\Phi^{-1}(a)} \ge c_4 \tag{7.7}$$

for some $c_4 > 0$. Using (7.3) and (7.7) in the second integral in (7.6), we get that for (a,r) with $\Phi(r) \le a \le (2^{-1} \wedge (2C_T)^{-2\delta_2})T$,

$$\begin{aligned} h_T(a,r) &\asymp a + \frac{a}{\Phi^{-1}(a)} + \int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2} ds + a \int_{\Phi^{-1}(a)}^{\Phi^{-1}(T)} \frac{1}{s^2} ds \\ &= a + \frac{a}{\Phi^{-1}(a)} + \int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2} ds + a \left(\frac{1}{\Phi^{-1}(a)} - \frac{1}{\Phi^{-1}(T)} \right) \\ &\asymp \frac{a}{\Phi^{-1}(a)} + \int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2} ds. \end{aligned} \tag{7.8}$$

Since $\Phi(s)$ is an increasing function, when $0 < \Phi(r) \le a$, we have

$$\frac{a}{\Phi^{-1}(a)} + \int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2} ds \le \frac{a}{\Phi^{-1}(a)} + a \int_r^{\Phi^{-1}(a)} \frac{1}{s^2} ds = \frac{a}{r},$$

while when $\Phi(r) \ge a > 0$,

$$\frac{a}{\Phi^{-1}(a)} + \left(\int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2} ds \right)^+ = \frac{a}{\Phi^{-1}(a)} \ge \frac{a}{r}.$$

This combined with (7.5) and (7.8) establishes the lemma. □

Recall that the Green function $G_D(x,y)$ of X in D is defined as $G_D(x,y) = \int_0^\infty p_D(t,x,y) dt$.

Theorem 7.3 *Suppose that X is a subordinate Brownian motion with Lévy exponent $\Psi(\xi) = \phi(|\xi|^2)$ with ϕ being a complete Bernstein function satisfying condition* **(A)**. *Let D be a bounded $C^{1,1}$ open subset of $\mathbb{R}^d$ with characteristics (R_2, Λ) and $a(x,y) = \Phi(\delta_D(x))^{1/2}\Phi(\delta_D(y))^{1/2}$, $x, y \in D$.*

(i) *There exists $c_1 > 0$ depending only on diam$(D), R_2, \Lambda, d$ and ϕ such that for all $d \geq 1$ and $(x,y) \in D \times D$,*

$$G_D(x,y) \geq c_1 \frac{\Phi(|x-y|)}{|x-y|^d}\left(1 \wedge \frac{\Phi(\delta_D(x))}{\Phi(|x-y|)}\right)^{1/2}\left(1 \wedge \frac{\Phi(\delta_D(y))}{\Phi(|x-y|)}\right)^{1/2}.$$

(ii) *There exists $c_2 > 0$ depending only on diam$(D), R_2, \Lambda, d$ and ϕ such that for all $d \geq 1$ and $(x,y) \in D \times D$,*

$$G_D(x,y) \leq c_2 \frac{a(x,y)}{|x-y|^d}.$$

(iii) *Let $d = 1$. Then for $(x,y) \in D \times D$,*

$$G_D(x,y) \asymp \frac{a(x,y)}{|x-y|} \wedge \left(\frac{a(x,y)}{\Phi^{-1}(a(x,y))} + \left(\int_{|x-y|}^{\Phi^{-1}(a(x,y))} \frac{\Phi(s)}{s^2}ds\right)^{+}\right).$$

(iv) *Let $d \geq 2$. Then for $(x,y) \in D \times D$,*

$$\begin{aligned} G_D(x,y) &\asymp \frac{\Phi(|x-y|)}{|x-y|^d}\left(1 \wedge \frac{\Phi(\delta_D(x))}{\Phi(|x-y|)}\right)^{1/2}\left(1 \wedge \frac{\Phi(\delta_D(y))}{\Phi(|x-y|)}\right)^{1/2} \\ &\asymp \frac{\Phi(|x-y|)}{|x-y|^d}\left(1 \wedge \frac{a(x,y)}{\Phi(|x-y|)}\right). \end{aligned}$$

Proof. Put $T = (2 \vee (2C_T)^{2\delta_2})\Phi(\mathrm{diam}(D))$, where C_T is the constant in (7.2). It follows from Theorem 1.5(iii) that

$$\int_T^\infty p_D(t,x,y)dt \asymp a(x,y). \tag{7.9}$$

Using the boundedness of D, Remark 1.1(iii) and (2.2), the results of Theorem 1.5(i)–(ii) can be rewritten as follows: there exists $c_1 > 0$ such that for $(t,x,y) \in (0,T] \times D \times D$,

$$\begin{aligned} & c_1^{-1}\left(1 \wedge \frac{\Phi(\delta_D(x))}{t}\right)^{1/2}\left(1 \wedge \frac{\Phi(\delta_D(y))}{t}\right)^{1/2}\left((\Phi^{-1}(t))^{-d} \wedge \frac{t}{|x-y|^d\Phi(|x-y|)}\right) \\ \leq\ & p_D(t,x,y) \qquad (7.10) \\ \leq\ & c_1\left(1 \wedge \frac{\Phi(\delta_D(x))}{t}\right)^{1/2}\left(1 \wedge \frac{\Phi(\delta_D(y))}{t}\right)^{1/2}\left((\Phi^{-1}(t))^{-d} \wedge \frac{t}{|x-y|^d\Phi(|x-y|)}\right). \qquad (7.11) \end{aligned}$$

By the change of variable $u = \frac{\Phi(|x-y|)}{t}$ and the fact that $t \to \Phi^{-1}(t)$ is increasing, we have

$$\begin{aligned}
&\int_0^T \left(1 \wedge \frac{\Phi(\delta_D(x))}{t}\right)^{1/2} \left(1 \wedge \frac{\Phi(\delta_D(y))}{t}\right)^{1/2} \left((\Phi^{-1}(t))^{-d} \wedge \frac{t}{|x-y|^d \Phi(|x-y|)}\right) dt \\
= \quad & \frac{\Phi(|x-y|)}{|x-y|^d} \left(\int_{\Phi(|x-y|)/T}^1 + \int_1^\infty\right) u^{-2} \left(\left(\frac{\Phi^{-1}(ut)}{\Phi^{-1}(t)}\right)^d \wedge u^{-1}\right) \left(1 \wedge \frac{\sqrt{u}\Phi(\delta_D(x))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) \\
& \qquad\qquad \times \left(1 \wedge \frac{\sqrt{u}\Phi(\delta_D(y))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) du \\
\asymp \quad & \frac{\Phi(|x-y|)}{|x-y|^d} \int_{\Phi(|x-y|)/T}^1 u^{-2} \left(\frac{|x-y|}{\Phi^{-1}(u^{-1}\Phi(|x-y|))}\right)^d \left(1 \wedge \frac{u a(x,y)}{\Phi(|x-y|)}\right) du \\
& + \frac{\Phi(|x-y|)}{|x-y|^d} \int_1^\infty u^{-3} \left(1 \wedge \frac{\sqrt{u}\Phi(\delta_D(x))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) \left(1 \wedge \frac{\sqrt{u}\Phi(\delta_D(y))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) du \\
=: \quad & I + II, \qquad (7.12)
\end{aligned}$$

where in the fourth line of the display above, we used Lemma 7.1.

(i) The estimate on II is easy.

$$\begin{aligned}
& \frac{1}{2} \frac{\Phi(|x-y|)}{|x-y|^d} \left(1 \wedge \frac{\Phi(\delta_D(x))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) \left(1 \wedge \frac{\Phi(\delta_D(y))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) \\
&= \frac{\Phi(|x-y|)}{|x-y|^d} \int_1^\infty u^{-3} \left(1 \wedge \frac{\Phi(\delta_D(x))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) \left(1 \wedge \frac{\Phi(\delta_D(y))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) du \\
\leq II \ &\leq \frac{\Phi(|x-y|)}{|x-y|^d} \int_1^\infty u^{-2} \left(u^{-1/2} \wedge \frac{\Phi(\delta_D(x))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) \left(u^{-1/2} \wedge \frac{\Phi(\delta_D(y))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) du \\
&\leq \frac{\Phi(|x-y|)}{|x-y|^d} \int_1^\infty u^{-2} \left(1 \wedge \frac{\Phi(\delta_D(x))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) \left(1 \wedge \frac{\Phi(\delta_D(y))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) du \\
&= \frac{\Phi(|x-y|)}{|x-y|^d} \left(1 \wedge \frac{\Phi(\delta_D(x))^{1/2}}{\Phi(|x-y|)^{1/2}}\right) \left(1 \wedge \frac{\Phi(\delta_D(y))^{1/2}}{\Phi(|x-y|)^{1/2}}\right). \qquad (7.13)
\end{aligned}$$

Now part (i) of the theorem follows from (7.10) and the lower bound of II in (7.13).

(ii) We let

$$u_0 := \frac{a(x,y)}{\Phi(|x-y|)}. \qquad (7.14)$$

Clearly $1/u_0 \geq \Phi(|x-y|)/\Phi(\mathrm{diam}(D)) \geq 2\Phi(|x-y|)/T$. By (7.2),

$$\begin{aligned}
I \quad &\leq \quad \frac{a(x,y)}{|x-y|^d} \int_{\Phi(|x-y|)/T}^1 \frac{|x-y|^d}{\Phi^{-1}(u^{-1}\Phi(|x-y|))^d} u^{-1} du \\
&= \quad \frac{a(x,y)}{|x-y|^d} \int_{\Phi(|x-y|)/T}^1 \left(\frac{\Phi^{-1}(\Phi(|x-y|))}{\Phi^{-1}(u^{-1}\Phi(|x-y|))}\right)^d u^{-1} du \\
&\leq \quad c_3 \frac{a(x,y)}{|x-y|^d} \int_0^1 u^{\frac{d}{2\delta_2}-1} du \\
&\leq \quad c_4 \frac{a(x,y)}{|x-y|^d}. \qquad (7.15)
\end{aligned}$$

Combining (7.9), (7.11), (7.13) and (7.15) we immediately get part (ii) of the theorem.

(iii) Let $h_T(a, r)$ be defined as in (7.4). Since $a \le \Phi(\mathrm{diam}(D)) \le (2^{-1} \wedge (2C_T)^{-2\delta_2}))T$, we have by (7.9)–(7.13) and Lemma 7.1 that $G_D(x, y) \asymp h_T(a(x, y), |x - y|)$. The assertion then follows from Lemma 7.2.

(iv) Note that since $d \ge 2$, we have by (7.2) that

$$\begin{aligned} I &= \frac{\Phi(|x-y|)}{|x-y|^d} \int_{\Phi(|x-y|)/T}^1 u^{-2} \left(\frac{|x-y|}{\Phi^{-1}(u^{-1}\Phi(|x-y|))} \right)^d \left(1 \wedge \frac{ua(x,y)}{\Phi(|x-y|)} \right) du \\ &\le c_5 \frac{\Phi(|x-y|)}{|x-y|^d} \left(1 \wedge \frac{a(x,y)}{\Phi(|x-y|)} \right) \int_0^1 u^{d/(2\delta_2)-2}\, du \\ &\le c_6 \frac{\Phi(|x-y|)}{|x-y|^d} \left(1 \wedge \frac{a(x,y)}{\Phi(|x-y|)} \right). \qquad (7.16) \end{aligned}$$

Part (iv) of the theorem now follows from assertion (i) of the theorem, Lemma 7.1, (7.9), (7.11), (7.13) and (7.16). □

Corollary 7.4 *Suppose that X is a one-dimensional subordinate Brownian motion with Lévy exponent $\Psi(\xi) = \phi(|\xi|^2)$ with ϕ being a complete Bernstein function satisfying condition* **(A)**. *Let D be a bounded $C^{1,1}$ open subset of $\mathbb{R}$ with characteristics (R_2, Λ) and $a(x, y) = \Phi(\delta_D(x))^{1/2}\Phi(\delta_D(y))^{1/2}$, $x, y \in D$.*

(i) *Suppose that for each $T > 0$ there is a constant $c_1 = c_1(T, \phi) > 0$ such that*

$$\int_r^T \frac{\Phi(s)}{s^2}\, ds \le c_1 \frac{\Phi(r)}{r} \qquad \text{for } r \in (0, T]. \qquad (7.17)$$

Then

$$G_D(x, y) \asymp \frac{\Phi(|x-y|)}{|x-y|} \left(1 \wedge \frac{a(x,y)}{\Phi(|x-y|)} \right).$$

(ii) *Suppose that for every $T > 0$, there is a constant $c_2 = c_2(T, \phi) > 0$ so that*

$$\int_0^r \frac{\Phi(s)}{s^2} ds \le c_2 \frac{\Phi(r)}{r} \qquad \text{for every } r \in (0, T]. \qquad (7.18)$$

Then for all $(x, y) \in D \times D$,

$$G_D(x, y) \asymp \frac{a(x,y)}{\Phi^{-1}(a(x,y))} \wedge \frac{a(x,y)}{|x-y|}.$$

Remark 7.5 Recall that $\delta_1, \delta_2 \in (0, 1)$ are the constants in condition **(A)**.

(i) Condition (7.17) is satisfied when $\delta_2 < 1/2$. This is because for $t \in (0, T]$, we have by (1.7) and condition **(A)**,

$$\frac{1}{\Phi(r)} \int_r^T \frac{\Phi(s)}{s^2}\, ds = \int_r^T \frac{\phi(r^{-2})}{\phi(s^{-2})} \frac{1}{s^2} ds \le a_2 \int_r^T \left(\frac{s^2}{r^2} \right)^{\delta_2} \frac{1}{s^2} ds = a_2 r^{-2\delta_2} \int_r^T s^{2\delta_2 - 2} ds \le \frac{c}{r}.$$

(ii) Condition (7.18) is satisfied when $\delta_1 > 1/2$. This is because for $t \in (0, T]$, we have by (1.7) and condition **(A)**,

$$\frac{1}{\Phi(r)}\int_0^r \frac{\Phi(s)}{s^2}\,ds = \int_0^r \frac{\phi(r^{-2})}{\phi(s^{-2})}\frac{1}{s^2}ds \le a_1\int_0^r \left(\frac{s^2}{r^2}\right)^{\delta_1}\frac{1}{s^2}ds = a_1 r^{-2\delta_2}\int_0^r s^{2\delta_1-2}ds \le \frac{c}{r}.$$

Remark 7.6 Let $\varphi(r) = r^{1/2}/\phi(r)$. Note that

$$a_1^{-1}\lambda^{1/2-\delta_1} \ge \varphi(\lambda r)/\varphi(r) \ge a_2^{-1}\lambda^{1/2-\delta_2} \qquad \text{for } \lambda \ge 1 \text{ and } r \ge R_0.$$

Let

$$\varphi^*(\lambda) := \limsup_{r\to\infty}\varphi(\lambda r)/\varphi(r) \quad \text{and} \quad \varphi_*(\lambda) := \liminf_{r\to\infty}\varphi(\lambda r)/\varphi(r).$$

Then

$$\infty > a_1^{-1}\lambda^{1/2-\delta_1} \ge \varphi^*(\lambda) \ge \varphi_*(\lambda) \ge a_2^{-1}\lambda^{1/2-\delta_2} > 0 \quad \text{for } \lambda \ge 1.$$

The upper and lower Matuszewska indices can be computed as

$$\alpha(\varphi) := \lim_{\lambda\to\infty}(\log\varphi_*(\lambda))/(\log\lambda), \quad \beta(\varphi) := \lim_{\lambda\to\infty}(\log\varphi^*(\lambda))/(\log\lambda).$$

Thus we have

$$\infty > a_1^{-1}\lambda^{1/2-\delta_1} \ge \varphi^*(\lambda) \ge \lambda^{\beta(\varphi)} \ge \lambda^{\alpha(\varphi)} \ge \varphi_*(\lambda) \ge a_2^{-1}\lambda^{1/2-\delta_2} > 0 \quad \text{for } \lambda \ge 1$$

(see [3, page 69-71]). Let $\widetilde{\varphi}(r) = \int_{T^{-1/2}}^r \varphi(t)/t\,dt$, $r \ge T^{-2}$. With the change of variable $s = t^{-1/2}$, we see that (7.17) is equivalent to

$$\widetilde{\varphi}(r) = \int_{T^{-2}}^r \frac{1}{\phi(t)t^{1/2}}dt = 2\int_{r^{-1/2}}^T \frac{\Phi(s)}{s^2}\,ds \le c_1\,\Phi(r^{-1/2})r^{1/2} = \varphi(r) \qquad \text{for } r \ge T^{-2}. \tag{7.19}$$

Let $\widehat{\varphi}(r) = \int_r^\infty \varphi(t)/t\,dt$. (7.18) is equivalent to

$$\widehat{\varphi}(r) = \int_r^\infty \frac{1}{\phi(t)t^{1/2}}dt = 2\int_0^{r^{-1/2}} \frac{\Phi(s)}{s^2}ds \le c_2\Phi(r^{-1/2})r^{1/2} = \varphi(r) \qquad \text{for every } r \ge T^{-2}. \tag{7.20}$$

In fact, by [3, Corollaries 2.6.2 and 2.6.4], $\varphi(r) \asymp \widetilde{\varphi}(r)$ for every $r \ge T^{-2}$ if and only if $\beta(\varphi) > 0$, and $\varphi(r) \asymp \widehat{\varphi}(r)$ for every $r \ge T^{-2}$ if and only if $\alpha(\varphi) < 0$.

By following the same proof in [21, Section 6], one can construct ϕ whose upper and lower Matuszewska indices are $\alpha(\phi) = 3/4$ and $\beta(\phi) = 1/4$ so that $\alpha(\varphi) = 1/4$ and $\beta(\varphi) = -1/4$. For such ϕ, neither (7.17) nor (7.18) hold.

Proof of Corollary 7.4. Put $T = (2 \vee (2C_T)^{2\delta_2})\Phi(\text{diam}(D))$, where $\text{diam}(D)$ is the diameter of D and C_T is the constant in (7.2). For notational simplicity, we let $a = a(x, y)$ and $r = |x - y|$. Recall that $u_0 = a/\Phi(r)$ is defined by (7.14). In view of Theorem 7.3(iii), we only need to consider $u_0 \ge 1$, which we will assume from now on in this proof.

(i) Since $a \le \Phi(\mathrm{diam}(D)) \le (2C_T)^{-2\delta_2}T$, by (7.7) and (7.17) we have

$$\begin{aligned}
&\frac{a}{\Phi^{-1}(a)} + \int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2}ds \asymp a\left(\frac{1}{\Phi^{-1}(a)} - \frac{1}{\Phi^{-1}(T)}\right) + \int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2}ds \\
&= a\int_{\Phi^{-1}(a)}^{\Phi^{-1}(T)} \frac{ds}{s^2} + \int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2}ds \le \int_r^{\Phi^{-1}(T)} \frac{\Phi(s)}{s^2}ds \le c_1\frac{\Phi(r)}{r}.
\end{aligned}$$

Combining this with Theorem 7.3(i), (iii) and Lemma 7.1 establishes part (i) of the corollary.

(ii) Since $a \le \Phi(\mathrm{diam}(D)) \le T/2$, by (7.18) we have that for $u_0 \ge 1$,

$$\frac{a}{\Phi^{-1}(a)} + \left(\int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2}ds\right)^+ \asymp \frac{a}{\Phi^{-1}(a)}.$$

Thus by Theorem 7.3(iii),

$$G_D(x,y) \asymp \frac{a}{\Phi^{-1}(a)} \qquad \text{when } u_0 \ge 1.$$

Since $\Phi(r)$ is increasing in r, the above together with Theorem 7.3(iii) implies that

$$G_D(x,y) \asymp \frac{a}{r} \wedge \frac{a}{\Phi^{-1}(a)} = \begin{cases} a/r & \text{if } a \le \Phi(r) \\ a/\Phi^{-1}(a) & \text{if } a \ge \Phi(r). \end{cases}$$

□

We next give an example of a one-dimensional subordinate Brownian motion with Lévy exponent $\phi(|\xi|^2)$ that satisfies condition **(A)** but its associated function $\Phi(r) = 1/\phi(r^{-2})$ satisfies neither condition (7.17) nor (7.18). We can get its explicit Green function estimates on bounded $C^{1,1}$ open sets by using Theorem 7.3(iii) but not by Corollary 7.4.

Example 7.7 By the discussion at the end of the Introduction, we know that for any $p \in \mathbb{R}$, the function

$$\phi(r) = \left(\int_0^\infty (r+t)^{-2}t^{1/2}(\ln(t+1))^p dt\right)^{-1}, \quad r > 0$$

is a complete Bernstein function. Moreover, $\phi(r) \asymp r^{1/2}(\log r)^p$ when $r \ge 2$. When $p \in [-1/2, 0) \cup (0, 1/2]$, $\phi(r) = r^{1/2}(\log(1+r))^p$ is an explicit example of a complete Bernstein function such that $\phi(r) \asymp r^{1/2}(\log r)^p$ when $r \ge 2$. Let X be a one-dimensional subordinate Brownian motion with Lévy exponent $\phi(|\xi|^2)$. Note that

$$\Phi(r) = 1/\phi(1/r^2) \asymp r(\log(1/r))^{-p} \quad \text{for } 0 < r \le 1/2. \tag{7.21}$$

Let $W(x)$ be the Lambert-W function, that is, $W(x)$ is the unique solution to $x = W(x)e^{W(x)}$. Suppose $s = r(\log(1/r))^{-p}$. Then

$$y := p^{-1}s^{-1/p} = r^{-1/p}\log((1/r)^{1/p}) =: xe^x.$$

So $\log r^{-1/p} = x = W(y) = W(p^{-1}s^{-1/p})$, which implies that $r = \exp(-pW(p^{-1}s^{-1/p}))$. So in view of (1.8), there is a constant $c_0 \in (0,1)$ so that

$$c_0 \exp(-pW(p^{-1}s^{-1/p})) \le \Phi^{-1}(s) \le c_0^{-1}\exp(-pW(p^{-1}s^{-1/p})) \qquad \text{for } s \in (0, \Phi(1/2)]. \tag{7.22}$$

Recall that

$$\int \frac{(\log(1/s))^{-p}}{s}ds = \frac{1}{p-1}(\log(1/s))^{1-p} + c, \qquad \text{when } p \neq 1$$

and

$$\int \frac{(\log(1/s))^{-1}}{s}ds = -\log(\log(1/s)) + c.$$

Suppose $0 < r \le \Phi^{-1}(a) \le 1/2$. Then if $p \neq 1$,

$$\begin{aligned}\int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2}ds &\asymp \int_r^{\Phi^{-1}(a)} \frac{(\log(1/s))^{-p}}{s}ds \\ &= \frac{1}{p-1}\left((\log(1/\Phi^{-1}(a)))^{1-p} - (\log(1/r))^{1-p}\right). \qquad (7.23)\end{aligned}$$

and, if $p = 1$,

$$\begin{aligned}\int_r^{\Phi^{-1}(a)} \frac{\Phi(s)}{s^2}ds &\asymp \int_r^{\Phi^{-1}(a)} \frac{(\log(1/s))^{-1}}{s}ds = \log\log(1/r) - \log\log(1/\Phi^{-1}(a)) \\ &= \log(\log r/\log\Phi^{-1}(a)). \qquad (7.24)\end{aligned}$$

Let D be a bounded $C^{1,1}$ open set in $\mathbb{R}$. We further assume that

$$\Phi^{-1}(\text{diam}(D)) \vee \text{diam}(D) < c_0/2,$$

where $c_0 \in (0,1)$ is the constant in (7.22).

Let $a(x,y) = \Phi(\delta_D(x))^{1/2}\Phi(\delta_D(y))^{1/2}$. We have by Theorem 7.3(iii), (7.21) and (7.23)-(7.24) that for $p \neq 1$,

$$\begin{aligned}G_D(x,y) &\asymp \frac{a(x,y)}{|x-y|} \wedge \left(\frac{a(x,y)}{\Phi^{-1}(a(x,y))} + \left(\int_{|x-y|}^{\Phi^{-1}(a(x,y))} \frac{\Phi(s)}{s^2}ds\right)^+\right) \\ &\asymp \frac{a(x,y)}{|x-y|} \wedge \left(\left(\log\frac{1}{\Phi^{-1}(a(x,y))}\right)^{-p} + \left(\frac{\left(\log(1/\Phi^{-1}(a(x,y)))\right)^{1-p} - (\log(1/|x-y|))^{1-p}}{p-1}\right)^+\right),\end{aligned}$$

while for $p = 1$,

$$G_D(x,y) \asymp \frac{a(x,y)}{|x-y|} \wedge \left(\left(\log\frac{1}{\Phi^{-1}(a(x,y))}\right)^{-1} + \log^+\left(\log|x-y|/\log\Phi^{-1}(a(x,y))\right)\right),$$

where $\log^+ x := 0 \vee \log x$. It is elementary to check that for $0 < u, r \le c_0/2$ and $c_0 u \le v \le c_0^{-1}u$,

$$\left(\log\frac{1}{u}\right)^{-p} + \left(\frac{(\log(1/u))^{1-p} - (\log(1/r))^{1-p}}{p-1}\right)^+ \asymp \left(\log\frac{1}{v}\right)^{-p} + \left(\frac{(\log(1/v))^{1-p} - (\log(1/r))^{1-p}}{p-1}\right)^+$$

when $p \neq 1$ and

$$\left(\log\frac{1}{u}\right)^{-1} + \log^+(\log r/\log u) \asymp \left(\log\frac{1}{v}\right)^{-1} + \log^+(\log r/\log v).$$

Thus we have from the four displays above together with (7.22) that for $p \neq 1$,

$$G_D(x,y) \asymp \frac{a(x,y)}{|x-y|} \wedge \left(W(p^{-1}a(x,y)^{1/p})^{-p} + \left(\frac{(W(p^{-1}a(x,y)^{1/p}))^{1-p} - (\log(1/|x-y|))^{1-p}}{p-1} \right)^+ \right),$$

while for $p = 1$,

$$G_D(x,y) \asymp \frac{a(x,y)}{|x-y|} \wedge \Big(W(a(x,y))^{-1} + \log^+ \big(W(a(x,y))^{-1} \log(1/|x-y|) \big) \Big).$$

Acknowledgement. The main results of this paper, in particular Theorems 1.3 and Corollary 1.6, were reported at the *The Sixth International Conference on Stochastic Analysis and Its Applications* held at Bedlewo, Poland, from September 9 to 14, 2012. At the same meeting, K. Bogdan, T. Grzywny and M. Ryznar announced that they had also obtained the factorization form estimates in terms of surviving probabilities and global heat kernel $p(t, c|x-y|)$ as in our Theorem 1.3(i)(ii) for the Dirichlet heat kernels of a similar class of purely discontinuous subordinate Brownian motions considered in this paper but only for bounded $C^{1,1}$ open sets.

References

[1] M. T. Barlow, R. F. Bass, Z.-Q. Chen and M. Kassmann. Non-local Dirichlet forms and symmetric jump processes. *Trans. Amer. Math. Soc.* **361** (2009), 1963–1999.

[2] M. T. Barlow, A. Grigor'yan and T. Kumagai. Heat kernel upper bounds for jump processes and the first exit time. *J. Reine Angew. Math.* **626** (2009), 135–157.

[3] N. H. Bingham, C. M. Goldie and J. L. Teugels, *Regular Variation*, Cambridge University Press, Cambridge, 1987.

[4] K. Bogdan, T. Grzywny and M. Ryznar. Heat kernel estimates for the fractional Laplacian with Dirichlet conditions. *Ann. Probab.* **38** (2010), 1901–1923.

[5] L. A. Caffarelli and L. Silvestre. Regularity theory for fully nonlinear integro-differential equations. *Comm. Pure Appl. Math.* **62(5)** (2009), 597–638.

[6] L. A. Caffarelli and L. Silvestre. The Evans-Krylov theorem for nonlocal fully nonlinear equations. *Ann. of Math. (2)* **174(2)** (2011), 1163–1187.

[7] L. A. Caffarelli, S. Salsa and L. Silvestre. Regularity estimates for the solution and the free boundary to the obstacle problem for the fractional Laplacian. *Invent. Math.* **171(1)** (2008) 425–461.

[8] Z.-Q. Chen, P. Kim and T. Kumagai. Weighted Poincaré Inequality and Heat Kernel Estimates for Finite Range Jump Processes. *Math. Ann.* **342(4)** (2008), 833–883.

[9] Z.-Q. Chen, P. Kim and T. Kumagai. On heat kernel estimates and parabolic Harnack inequality for jump processes on metric measure spaces. *Acta Mathematica Sinica, English Series* **25** (2009), 1067–1086.

[10] Z.-Q. Chen, P. Kim and T. Kumagai. Global Heat Kernel Estimates for Symmetric Jump Processes. *Trans. Amer. Math. Soc.* **363(9)** (2012), 5021–5055.

[11] Z.-Q. Chen, P. Kim, and R. Song. Heat kernel estimates for Dirichlet fractional Laplacian. *J. European Math. Soc.*, **12** (2010), 1307–1329.

[12] Z.-Q. Chen, P. Kim and R. Song. Two-sided heat kernel estimates for censored stable-like processes. *Probab. Theory Relat. Fields*, **146** (2010), 361–399.

[13] Z.-Q. Chen, P. Kim and R. Song. Dirichlet heat kernel estimates for $\Delta^{\alpha/2}+\Delta^{\beta/2}$. *Ill. J. Math.* **54** (2010) (Special issue in honor of D. Burkholder), 1357–1392.

[14] Z.-Q. Chen, P. Kim and R. Song. Sharp heat kernel estimates for relativistic stable processes in open sets. *Ann. Probab.* **40** (2012), 213–244.

[15] Z.-Q. Chen, P. Kim and R. Song. Heat kernel estimate for $\Delta + \Delta^{\alpha/2}$ in $C^{1,1}$ open sets. *J. London Math. Soc.* **84 (1)** (2011), 58–80.

[16] Z.-Q. Chen, P. Kim, and R. Song. Global heat kernel estimates for relativistic stable processes in half-space-like open sets. *Potential Anal.*, **36** (2012) 235–261.

[17] Z.-Q. Chen and T. Kumagai. Heat kernel estimates for stable-like processes on d-sets. *Stoch. Proc. Appl.* **108** (2003), 27–62.

[18] Z.-Q. Chen and T. Kumagai. Heat kernel estimates for jump processes of mixed types on metric measure spaces. *Probab. Theory Relat. Fields*, **140** (2008), 277–317.

[19] P. Kim and A. Mimica. Green function estimates for subordinate Brownian motions: stable and beyond. To appear in *Trans. Amer. Math. Soc.*

[20] P. Kim, R. Song and Z. Vondraček. Two-sided Green function estimates for killed subordinate Brownian motions. *Proc. London Math. Soc.* **104** (2012), 927–958.

[21] P. Kim, R. Song and Z. Vondracek. Uniform boundary Harnack principle for rotationally symmetric Lévy processes in general open sets. *Sci. China Math.* **55**, (2012), 2193–2416.

[22] P.-A. Meyer. Renaissance, recollements, mélanges, ralentissement de processus de Markov. *Ann. Inst. Fourier*, **25** (1975), 464–497.

[23] R. L. Schilling, R. Song and Z. Vondraček. *Bernstein Functions: Theory and Applications.* de Gruyter Studies in Mathematics 37. Berlin: Walter de Gruyter, 2010.

[24] L. Silvestre. Hölder estimates for solutions of integro differential equations like the fractional Laplacian. *Indiana Univ. Math. J.* **55** (2006), 1155–1174.

[25] T. Watanabe. The isoperimetric inequality for isotropic unimodal Lévy processes. *Z. Wahrs. verw. Gebiete* **63** (1983), 487–499.

[26] S. Watanabe, K. Yano and Y. Yano. A density formula for the law of time spent on the positive side of one-dimensional diffusion processes. *J. Math. Kyoto Univ.* **45** (2005), 781–806.

[27] M. Zähle. Potential spaces and traces of Lévy processes on h-sets. *J. Contemp. Math. Anal.* **44** (2009), 117–145.

Zhen-Qing Chen
Department of Mathematics, University of Washington, Seattle, WA 98195, USA
E-mail: `zqchen@uw.edu`

Panki Kim
Department of Mathematics, Seoul National University, Seoul 151-742, South Korea
E-mail: `pkim@snu.ac.kr`

Renming Song
Department of Mathematics, University of Illinois, Urbana, IL 61801, USA
E-mail: `rsong@math.uiuc.edu`